

Math 1553 Worksheet §3.3 and §3.4, Matrix Equations and Solution Sets
Solutions

For problems 1 and 2 below: The professor in your widgets and gizmos class is trying to decide between three different grading schemes for computing your final course grade. The schemes are based on homework (HW), quiz grades (Q), midterms (M), and a final exam (F). The three schemes can be described by the following matrix A:

$$\begin{array}{r} \text{Scheme 1} \\ \text{Scheme 2} \\ \text{Scheme 3} \end{array} \begin{array}{cccc} \text{HW} & \text{Q} & \text{M} & \text{F} \\ \left(\begin{array}{cccc} 0.1 & 0.1 & 0.5 & 0.3 \\ 0.1 & 0.1 & 0.4 & 0.4 \\ 0.1 & 0.1 & 0.6 & 0.2 \end{array} \right) \end{array}$$

Feel free to use a calculator to carry out arithmetic in problems 1 and 2.

1. Suppose that you have a score of x_1 on homework, x_2 on quizzes, x_3 on midterms, and x_4 on the final, with potential final course grades of b_1, b_2, b_3 . Write a matrix equation $Ax = b$ to relate your final grades to your scores.

Solution.

In the above grading schemes, you would receive the following final grades:

$$\begin{array}{l} \text{Scheme 1: } 0.1x_1 + 0.1x_2 + 0.5x_3 + 0.3x_4 = b_1 \\ \text{Scheme 2: } 0.1x_1 + 0.1x_2 + 0.4x_3 + 0.4x_4 = b_2 \\ \text{Scheme 3: } 0.1x_1 + 0.1x_2 + 0.6x_3 + 0.2x_4 = b_3 \end{array}$$

This is the same as the matrix equation

$$(*) \quad \begin{pmatrix} 0.1 & 0.1 & 0.5 & 0.3 \\ 0.1 & 0.1 & 0.4 & 0.4 \\ 0.1 & 0.1 & 0.6 & 0.2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}.$$

2. Suppose that you end up with averages of 90% on the homework, 90% on quizzes, 85% on midterms, and a 95% score on the final exam. Use Problem 1 to determine which grading scheme leaves you with the highest overall course grade.

Solution.

According to equation (*) above, your final grades would be

$$\begin{pmatrix} 0.1 & 0.1 & 0.5 & 0.3 \\ 0.1 & 0.1 & 0.4 & 0.4 \\ 0.1 & 0.1 & 0.6 & 0.2 \end{pmatrix} \begin{pmatrix} .90 \\ .90 \\ .85 \\ .95 \end{pmatrix} = \begin{pmatrix} .89 \\ .90 \\ .88 \end{pmatrix}.$$

Hence the second grading scheme gives you the best final grade.

3. If the statement is always true, circle TRUE. Otherwise, circle FALSE. Justify your answer.

a) If A is a 5×4 matrix, then the equation $Ax = b$ must be inconsistent for some b in \mathbf{R}^5 .

TRUE FALSE

b) Any linear combination of vectors can always be written in the form Ax for a suitable matrix A and vector x .

TRUE FALSE

Solution.

a) True. If A is a 5×4 matrix, then A can have at most 4 pivots (since no row or column can have more than 1 pivot). But A has 5 rows, so this means A cannot have a pivot in each row, and therefore $Ax = b$ must be inconsistent for at least one b in \mathbf{R}^5 .

b) True. Any linear combination of vectors in \mathbf{R}^n (any n) can be written

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p$$

for some scalars x_1, \dots, x_p and vectors v_1, \dots, v_p in \mathbf{R}^n . This linear combination is Ax where

$$A = (v_1 \quad v_2 \quad \cdots \quad v_p) \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{pmatrix}.$$

4. Find the solution sets of $x_1 - 3x_2 + 5x_3 = 0$ and $x_1 - 3x_2 + 5x_3 = 3$ and write them in parametric vector form. How do the solution sets compare geometrically?

Solution.

The equation $x_1 - 3x_2 + 5x_3 = 0$ corresponds to the augmented matrix $(1 \quad -3 \quad 5 \mid 0)$ which has two free variables x_2 and x_3 .

$$x_1 = 3x_2 - 5x_3 \quad x_2 = x_2 \quad x_3 = x_3.$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3x_2 - 5x_3 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3x_2 \\ x_2 \\ 0 \end{pmatrix} + \begin{pmatrix} -5x_3 \\ 0 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -5 \\ 0 \\ 1 \end{pmatrix}.$$

The solution set for $x_1 - 3x_2 + 5x_3 = 0$ is the plane spanned by $\begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} -5 \\ 0 \\ 1 \end{pmatrix}$.

The equation $x_1 - 3x_2 + 5x_3 = 3$ corresponds to the augmented matrix $(1 \quad -3 \quad 5 \mid 3)$ which has two free variables x_2 and x_3 .

$$x_1 = 3 + 3x_2 - 5x_3 \quad x_2 = x_2 \quad x_3 = x_3.$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 + 3x_2 - 5x_3 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 3x_2 \\ x_2 \\ 0 \end{pmatrix} + \begin{pmatrix} -5x_3 \\ 0 \\ x_3 \end{pmatrix} = \boxed{\begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -5 \\ 0 \\ 1 \end{pmatrix}}.$$

This solution set is the *translation* by $\begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}$ of the plane spanned by $\begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} -5 \\ 0 \\ 1 \end{pmatrix}$.