

Supplemental problems: §5.4

1. True or false. Answer true if the statement is always true. Otherwise, answer false.
 - a) If A is an invertible matrix and A is diagonalizable, then A^{-1} is diagonalizable.
 - b) A diagonalizable $n \times n$ matrix admits n linearly independent eigenvectors.
 - c) If A is diagonalizable, then A has n distinct eigenvalues.
2. Give examples of 2×2 matrices with the following properties. Justify your answers.
 - a) A matrix A which is invertible and diagonalizable.
 - b) A matrix B which is invertible but not diagonalizable.
 - c) A matrix C which is not invertible but is diagonalizable.
 - d) A matrix D which is neither invertible nor diagonalizable.

3. $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}.$


- a) Find the eigenvalues of A , and find a basis for each eigenspace.
- b) Is A diagonalizable? If your answer is yes, find a diagonal matrix D and an invertible matrix C so that $A = CDC^{-1}$. If your answer is no, justify why A is not diagonalizable.

4. Let $A = \begin{pmatrix} 8 & 36 & 62 \\ -6 & -34 & -62 \\ 3 & 18 & 33 \end{pmatrix}.$

The characteristic polynomial for A is $-\lambda^3 + 7\lambda^2 - 16\lambda + 12$, and $\lambda - 3$ is a factor. Decide if A is diagonalizable. If it is, find an invertible matrix C and a diagonal matrix D such that $A = CDC^{-1}$.

5. Which of the following 3×3 matrices are necessarily diagonalizable over the real numbers? (Circle all that apply.)
 1. A matrix with three distinct real eigenvalues.
 2. A matrix with one real eigenvalue.
 3. A matrix with a real eigenvalue λ of algebraic multiplicity 2, such that the λ -eigenspace has dimension 2.
 4. A matrix with a real eigenvalue λ such that the λ -eigenspace has dimension 2.

6. Suppose a 2×2 matrix A has eigenvalue $\lambda_1 = -2$ with eigenvector $v_1 = \begin{pmatrix} 3/2 \\ 1 \end{pmatrix}$, and eigenvalue $\lambda_2 = -1$ with eigenvector $v_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$.
- Find A .
 - Find A^{100} .
7. Suppose that $A = C \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix} C^{-1}$, where C has columns v_1 and v_2 . Given x and y in the picture below, draw the vectors Ax and Ay .



Supplemental problems: §5.5

1.
 - a) If A is the matrix that implements rotation by 143° in \mathbf{R}^2 , then A has no real eigenvalues.
 - b) A 3×3 matrix can have eigenvalues $3, 5$, and $2 + i$.
 - c) If $v = \begin{pmatrix} 2+i \\ 1 \end{pmatrix}$ is an eigenvector of A corresponding to the eigenvalue $\lambda = 1 - i$, then $w = \begin{pmatrix} 2i-1 \\ i \end{pmatrix}$ is an eigenvector of A corresponding to the eigenvalue $\lambda = 1 - i$.

2. Consider the matrix

$$A = \begin{pmatrix} 3\sqrt{3}-1 & -5\sqrt{3} \\ 2\sqrt{3} & -3\sqrt{3}-1 \end{pmatrix}$$

- a) Find both complex eigenvalues of A .
 - b) Find an eigenvector corresponding to each eigenvalue.
3. Let $A = \begin{pmatrix} 4 & -3 & 3 \\ 3 & 4 & -2 \\ 0 & 0 & 2 \end{pmatrix}$. Find all eigenvalues of A . For each eigenvalue of A , find a corresponding eigenvector.