

Math 1553 Worksheet §§3.4-3.6

Solutions

1. True or false (justify your answer). Answer true if the statement is *always* true. Otherwise, answer false.
- a) If A is a 3×4 matrix and B is a 4×2 matrix, then the linear transformation Z defined by $Z(x) = ABx$ has domain \mathbf{R}^3 and codomain \mathbf{R}^2 .
 - b) If A is an $n \times n$ matrix and the equation $Ax = b$ has at least one solution for each b in \mathbf{R}^n , then the solution is *unique* for each b in \mathbf{R}^n .
 - c) Suppose A is an $n \times n$ matrix and every vector in \mathbf{R}^n can be written as a linear combination of the columns of A . Then A must be invertible.

Solution.

- a) False. In order for Bx to make sense, x must be in \mathbf{R}^2 , and so Bx is in \mathbf{R}^4 and $A(Bx)$ is in \mathbf{R}^3 . Therefore, the domain of Z is \mathbf{R}^2 and the codomain of Z is \mathbf{R}^3 .
- b) True. The first part says the transformation $T(x) = Ax$ is onto. Since A is $n \times n$, this is the same as saying A is invertible, so T is one-to-one and onto. Therefore, the equation $Ax = b$ has exactly one solution for each b in \mathbf{R}^n .
- c) True. If the columns of A span \mathbf{R}^n , then A is invertible by the Invertible Matrix Theorem. We can also see this directly without quoting the IMT:
If the columns of A span \mathbf{R}^n , then A has n pivots, so A has a pivot in each row and column, hence its matrix transformation $T(x) = Ax$ is one-to-one and onto and thus invertible. Therefore, A is invertible.

2. Let $T : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be rotation *clockwise* by 60° . Let $U : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be the linear transformation satisfying $U(1, 0) = (-2, 1)$ and $U(0, 1) = (1, 0)$.
- Find the standard matrix for the composition $U \circ T$ using matrix multiplication.
 - Find the standard matrix for the composition $T \circ U$ using matrix multiplication.
 - Is rotating clockwise by 60° and then performing U , the same as first performing U and then rotating clockwise by 60° ?

Solution.

a) The matrix for T is $\begin{pmatrix} \cos(-60^\circ) & -\sin(-60^\circ) \\ \sin(-60^\circ) & \cos(-60^\circ) \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$.

The matrix for U is $(U(e_1) \ U(e_2)) = \begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix}$.

The matrix for $U \circ T$ is

$$\begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} = \begin{pmatrix} -1 - \frac{\sqrt{3}}{2} & \frac{1}{2} - \sqrt{3} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix}.$$

b) The matrix for $T \circ U$ is

$$\begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 + \frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{1}{2} + \sqrt{3} & -\frac{\sqrt{3}}{2} \end{pmatrix}.$$

- c) No. In (a) and (b), we found that the standard matrices for $U \circ T$ and $T \circ U$ are different, so the transformations are different.