MATH 1553, SPRING 2019 SAMPLE MIDTERM 3: 5.1 THROUGH 6.6

Name	ГEmail	@gatech.edu
------	--------	-------------

Write your section number here:

Please read all instructions carefully before beginning.

- The maximum score on this exam is 50 points.
- You have 50 minutes to complete this exam.
- There are no aids of any kind (calculators, notes, text, etc.) allowed.
- Please show your work unless instructed otherwise. A correct answer without appropriate work will receive little or no credit. If you cannot fit your work on the front side of the page, use the back side of the page as indicated.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 50 minutes, without notes or distractions.

The exam is not designed to test material from the previous midterm on its own. However, knowledge of the material prior to section §5.1 is necessary for everything we do for the rest of the semester, so it is fair game for the exam as it applies to §§5.1 through 6.6.


Answer true if the statement is <i>always</i> true. Otherwise, answer false. In every case, assume that the entries of the matrix <i>A</i> are real numbers.				
a)	Т	F	If <i>A</i> is the 3 × 3 matrix satisfying $Ae_1 = e_2$, $Ae_2 = e_3$, and $Ae_3 = e_1$, then det(<i>A</i>) = 1.	
b)	Т	F	If <i>A</i> is an $n \times n$ matrix and det(<i>A</i>) = 2, then 2 is an eigenvalue of <i>A</i> .	
c)	Т	F	If <i>A</i> and <i>B</i> are $n \times n$ matrices with det(<i>A</i>) = 0 and det(<i>B</i>) = 0, then det(<i>A</i> + <i>B</i>) = 0.	
d)	Т	F	If <i>A</i> is an $n \times n$ matrix and <i>v</i> and <i>w</i> are eigenvectors of <i>A</i> , then $v + w$ is also an eigenvector of <i>A</i> .	
e)	Т	F	If <i>A</i> is a positive 2 × 2 stochastic matrix with steady-state vector $v = \begin{pmatrix} 7/10 \\ 3/10 \end{pmatrix}$, then $A^n \begin{pmatrix} 40 \\ 60 \end{pmatrix}$ approaches $\begin{pmatrix} 70 \\ 30 \end{pmatrix}$ as $n \to \infty$.	

Extra space for scratch work on problem 1

Problem 2.

Short answer. Show your work on part (c). In every case, the entries of each matrix must be real numbers.

- a) Write a 2×2 matrix *A* which is invertible but not diagonalizable.
- **b)** Write a 2 × 2 matrix *A* for which $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are eigenvectors corresponding to the same eigenvalue.
- **c)** Find the area of the parallelogram drawn below (the grid marks are spaced one unit apart).



d) Write a 3 × 3 matrix *A* with only one real eigenvalue $\lambda = 4$, such that the 4-eigenspace for *A* is a two-dimensional plane in \mathbb{R}^3 .

e) Suppose *A* is an $n \times n$ matrix. Which of the following must be true? Circle all that apply.

I. If det(A) = 0 then *A* is not invertible.

II. If *A* is diagonalizable, then *A* has *n* distinct eigenvalues.

Problem 3.

Parts (a) and (b) are unrelated.

a) Consider the matrix

$$A = \begin{pmatrix} 3 & -7 \\ 1 & -1 \end{pmatrix}$$

Find all eigenvalues of *A*. Simplify your answer. For the eigenvalue with negative imaginary part, find an eigenvector.

b) Consider an internet with four pages: 1, 2, 3, and 4. Page 1 links to pages 2 and 3. Page 2 links to pages 3 and 4. Page 3 links to pages 1 and 4. Page 4 links to page 1.

(i) Write the importance matrix *A* for this internet.

(ii) *A* has a unique steady-state vector *w*. Find *w*. You may use the fact that $A \begin{pmatrix} 8 \\ 4 \\ 6 \\ 5 \end{pmatrix} = \begin{pmatrix} 8 \\ 4 \\ 6 \\ 5 \end{pmatrix}.$

Problem 4.

 $\operatorname{Let} A = \begin{pmatrix} -1 & 0 & -2 \\ 0 & 2 & 0 \\ 3 & 0 & 4 \end{pmatrix}.$ **a)** Find the eigenvalues of *A*. **b)** Find a basis for each eigenspace of *A*. Mark your answers clearly. c) Is A diagonalizable? If your answer is yes, find a diagonal matrix D and an invertible matrix C so that $A = CDC^{-1}$. If your answer is no, justify why A is not diagonalizable.

Problem 5.

[10 points]

Parts (a) and (b) are not related.

a) Find det(
$$A^3$$
) if $A = \begin{pmatrix} 1 & -3 & 4 & 2 \\ 0 & 0 & -2 & 0 \\ 0 & 1 & 2 & 3 \\ 2 & 0 & -1 & 20 \end{pmatrix}$.

b) Find the 2×2 matrix *A* whose eigenspaces are drawn below. Fully simplify your answer. (to be clear: the dashed line is the (-2)-eigenspace).

