

**MATH 1553, FALL 2018**  
**SAMPLE MIDTERM 1: THROUGH SECTION 3.4**

<b>Name</b>	
-------------	--

Please **read all instructions** carefully before beginning.

- You have 50 minutes to complete this exam.
- There are no aids of any kind (calculators, notes, text, etc.) allowed.
- Please show your work unless specified otherwise. A correct answer without appropriate work may be given little or no credit.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 50 minutes, without notes or distractions.

## Problem 1.

[Parts a) through f) are worth 2 points each]

a) Compute:  $\begin{pmatrix} 3 & 2 \\ -2 & 0 \\ 1 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ -3 \end{pmatrix} =$

The remaining problems are True or false. Circle **T** if the statement is **always** true, and circle **F** otherwise. You do not need to justify your answer.

b) **T**    **F**    The matrix  $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$  is in reduced row echelon form.

c) **T**    **F**    If  $Ax = b$  is consistent, then the equation  $Ax = 5b$  is consistent.

d) **T**    **F**    If the augmented matrix corresponding to a linear system of equations has a pivot in every row, then the system is consistent.

e) **T**    **F**    If  $A$  is an  $m \times n$  matrix and  $Ax = 0$  has a unique solution, then  $Ax = b$  is consistent for every  $b$  in  $\mathbf{R}^m$ .

f) **T**    **F**    The three vectors  $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ , and  $\begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$  span  $\mathbf{R}^3$ .

### Solution.

a)  $1 \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} - 3 \begin{pmatrix} 2 \\ 0 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} + \begin{pmatrix} -6 \\ 0 \\ -12 \end{pmatrix} = \begin{pmatrix} -3 \\ -2 \\ -11 \end{pmatrix}.$

b) True.

c) True. If  $Aw = b$  then  $A(5w) = 5Aw = 5b$ .

d) False. For example,  $\left( \begin{array}{cc|c} \boxed{1} & 0 & 0 \\ 0 & 0 & \boxed{1} \end{array} \right)$  has a pivot in every row but is inconsistent.

e) False. For example, if  $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$ , then  $Ax = 0$  has only the trivial solution, but

$$Ax = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \text{ has no solution.}$$

f) True. The three vectors form a  $3 \times 3$  matrix with a pivot in every row.

## Problem 2.

Parts (a) and (b) are 2 points each. Parts (c) and (d) are 3 points each.

a) If  $A$  is a  $2 \times 3$  matrix with 2 pivots, then the set of solutions to  $Ax = 0$  is a:


(circle one answer)    **point**    **line**    **plane**    **3-plane**

in:

(circle one answer)    **R**    **R<sup>2</sup>**    **R<sup>3</sup>**.

b) Write a vector equation which represents an inconsistent system of two linear equations in  $x_1$  and  $x_2$ .

c) For some  $2 \times 2$  matrix  $A$  and vector  $b$  in  $\mathbf{R}^2$ , the solution set of  $Ax = b$  is drawn below. Draw the solution set of  $Ax = 0$ .



d) If  $b, v, w$  are vectors in  $\mathbf{R}^3$  and  $\text{Span}\{b, v, w\} = \mathbf{R}^3$ , is it possible that  $b$  is in  $\text{Span}\{v, w\}$ ? Justify your answer.

### Solution.

a) Line in  $\mathbf{R}^3$ . Since there are 2 pivots but 3 columns, one column will not have a pivot, so  $Ax = 0$  will have exactly one free variable. The number of entries in  $x$  must match the number of columns of  $A$  (namely, 3), so each solution  $x$  is in  $\mathbf{R}^3$ .

b) The system  $\begin{pmatrix} 1 & 1 & | & 0 \\ 1 & 1 & | & 1 \end{pmatrix}$  is inconsistent; its corresponding vector equation is

$$x_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

c) The solution set of  $Ax = 0$  is the parallel line through the origin.

d) No. Recall that  $\text{Span}\{b, v, w\}$  is the set of all linear combinations of  $b, v$ , and  $w$ . If  $b$  is in  $\text{Span}\{v, w\}$  then  $b$  is a linear combination of  $v$  and  $w$ . Consequently, any element of  $\text{Span}\{b, v, w\}$  is a linear combination of  $v$  and  $w$  and is therefore in  $\text{Span}\{v, w\}$ ,

which is at most a plane and cannot be all of  $\mathbf{R}^3$ .

To see why the span of  $v$  and  $w$  can never be  $\mathbf{R}^3$ , consider the matrix  $A$  whose columns are  $v$  and  $w$ . Since  $A$  is  $3 \times 2$ , it has at most two pivots, so  $A$  cannot have a pivot in every row. Therefore, by a theorem from section 1.4, the equation  $Ax = b$  will fail to be consistent for some  $b$  in  $\mathbf{R}^3$ , which means that some  $b$  in  $\mathbf{R}^3$  is not in the span of  $v$  and  $w$ .

### Problem 3.

[10 points]

Johnny Rico believes that the secret to the universe can be found in the system of two linear equations in  $x$  and  $y$  given by

$$\begin{aligned}x - y &= h \\ 3x + hy &= 4\end{aligned}$$

where  $h$  is a real number.

- Find all values of  $h$  (if any) which make the system inconsistent. Briefly justify your answer.
- Find all values of  $h$  (if any) which make the system have a unique solution. Briefly justify your answer.

### Solution.

Represent the system with an augmented matrix and row-reduce:

$$\left( \begin{array}{cc|c} 1 & -1 & h \\ 3 & h & 4 \end{array} \right) \xrightarrow{R_2 - 3R_1} \left( \begin{array}{cc|c} 1 & -1 & h \\ 0 & h+3 & 4-3h \end{array} \right).$$

- If  $h = -3$  then the matrix is  $\left( \begin{array}{cc|c} 1 & -1 & -3 \\ 0 & 0 & 13 \end{array} \right)$ , which has a pivot in the rightmost column and is therefore inconsistent.
- If  $h \neq -3$ , then the matrix has a pivot in each row to the left of the augment:  $\left( \begin{array}{cc|c} \boxed{1} & -1 & h \\ 0 & \boxed{h+3} & 4-3h \end{array} \right)$ . The right column is not a pivot column, so the system is consistent. The left side has a pivot in each column, so the solution is unique.

## Problem 4.

[11 points]

- a) Find the parametric form of the general solution of the following system of equations. Clearly indicate which variables (if any) are free variables.

$$\begin{aligned}x_1 + 2x_2 + 2x_3 - x_4 &= 4 \\2x_1 + 4x_2 + x_3 - 2x_4 &= -1 \\-x_1 - 2x_2 - x_3 + x_4 &= -1\end{aligned}$$

- b) Write the set of solutions to

$$\begin{aligned}x_1 + 2x_2 + 2x_3 - x_4 &= 0 \\2x_1 + 4x_2 + x_3 - 2x_4 &= 0 \\-x_1 - 2x_2 - x_3 + x_4 &= 0\end{aligned}$$

in parametric vector form.

### Solution.

- a) We put the appropriate augmented matrix into RREF.

$$\begin{aligned}\left(\begin{array}{cccc|c}1 & 2 & 2 & -1 & 4 \\2 & 4 & 1 & -2 & -1 \\-1 & -2 & -1 & 1 & -1\end{array}\right) &\xrightarrow[\substack{R_2=R_2-2R_1 \\ R_3=R_3+R_1}]{R_2 \leftrightarrow R_3} \left(\begin{array}{cccc|c}1 & 2 & 2 & -1 & 4 \\0 & 0 & -3 & 0 & -9 \\0 & 0 & 1 & 0 & 3\end{array}\right) \xrightarrow{R_2 \leftrightarrow R_3} \left(\begin{array}{cccc|c}1 & 2 & 2 & -1 & 4 \\0 & 0 & 1 & 0 & 3 \\0 & 0 & -3 & 0 & -9\end{array}\right) \\ &\xrightarrow[\substack{R_3=R_3+3R_2 \\ R_1=R_1-2R_2}]{R_3=R_3+3R_2} \left(\begin{array}{cccc|c}1 & 2 & 0 & -1 & -2 \\0 & 0 & 1 & 0 & 3 \\0 & 0 & 0 & 0 & 0\end{array}\right)\end{aligned}$$

Therefore,  $x_2$  and  $x_4$  are free, and we have:

$$\begin{aligned}x_1 &= -2 - 2x_2 + x_4 \\x_2 &= x_2 \\x_3 &= 3 \\x_4 &= x_4.\end{aligned}$$

In parametric form, this is:

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -2 - 2x_2 + x_4 \\ x_2 \\ 3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \\ 3 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$


- b) The equation in (b) is just the corresponding homogeneous equation, which is a translate of the above plane which includes the origin.

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \quad (x_2, x_4 \text{ real}).$$

## Problem 5.

[7 points]

The diagram below represents traffic in a city.



- a) Write a system of three linear equations whose solution would give the values of  $x_1$ ,  $x_2$ , and  $x_3$ . Do not solve it.
- b) Write the system of equations as a vector equation. Do not solve it.

### Solution.

- a) The number of cars leaving an intersection must equal the number of cars entering.

$$x_3 + 70 = x_1 + 90$$

$$x_1 + x_2 = 160$$

$$x_2 + x_3 = 180.$$

Or:

$$-x_1 + x_3 = 20$$

$$x_1 + x_2 = 160$$

$$x_2 + x_3 = 180.$$

b)  $x_1 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + x_3 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 20 \\ 160 \\ 180 \end{pmatrix}.$

[Scratch work]