

### Supplemental problems: §2.1, §2.2

1. Consider the augmented matrix

$$\left( \begin{array}{ccc|c} 2 & -2 & 2 & 0 \\ 1 & -3 & -4 & -9 \\ 3 & -1 & 8 & 9 \end{array} \right)$$

**Question:** Does the corresponding linear system have a solution? If so, what is the solution set?

- Formulate this question as a vector equation.
- Formulate this question as a system of linear equations.
- What does this mean in terms of spans?
- Answer the question using the [interactive demo](#).
- Answer the question using row reduction.
- Find a **different** solution in parts (e) and (d).

**Solution.**

a) What are the solutions to the following vector equation?

$$x \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + y \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix} + z \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix} = \begin{pmatrix} 0 \\ -9 \\ 9 \end{pmatrix}$$

b) What is the solution set of the following linear system?

$$\begin{aligned} 2x - 2y + 2z &= 0 \\ x - 3y - 4z &= -9 \\ 3x - y + 8z &= 9 \end{aligned}$$

c) There exists a solution if and only if  $\begin{pmatrix} 0 \\ -9 \\ 9 \end{pmatrix}$  is in  $\text{Span} \left\{ \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix}, \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix} \right\}$ .

e) Row reducing yields

$$\left( \begin{array}{ccc|c} 1 & 0 & 7/2 & 9/2 \\ 0 & 1 & 5/2 & 9/2 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Hence  $z$  is a free variable, so the solution in parametric form is

$$\begin{aligned} x &= \frac{9}{2} - \frac{7}{2}z \\ y &= \frac{9}{2} - \frac{5}{2}z. \end{aligned}$$

Taking  $z = 0$  yields the solution  $x = y = 9/2$ .

f) Taking  $z = 1$  yields the solution  $x = 1, y = 2$ .

2. Let  $v_1 = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$      $v_2 = \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix}$      $w = \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix}$ .

**Question:** Is  $w$  a linear combination of  $v_1$  and  $v_2$ ? In other words, is  $w$  in  $\text{Span}\{v_1, v_2\}$ ?

- Formulate this question as a vector equation.
- Formulate this question as a system of linear equations.
- Formulate this question as an augmented matrix.
- Answer the question using the [interactive demo](#).
- Answer the question using row reduction.

**Solution.**

- a) Does the following vector equation have a solution?

$$x \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + y \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix}$$

- b) Does the following linear system have a solution?

$$\begin{aligned} 2x - 2y &= 2 \\ x - 3y &= -4 \\ 3x - y &= 8 \end{aligned}$$

- c) As an augmented matrix:

$$\left( \begin{array}{cc|c} 2 & -2 & 2 \\ 1 & -3 & -4 \\ 3 & -1 & 8 \end{array} \right)$$

- e) Row reducing yields

$$\left( \begin{array}{cc|c} 1 & 0 & 7/2 \\ 0 & 1 & 5/2 \\ 0 & 0 & 0 \end{array} \right)$$

so  $x = 7/2$  and  $y = 5/2$ .

3. Let

$$A = \begin{pmatrix} 1 & 0 & 5 \\ -2 & 1 & -6 \\ 0 & 2 & 8 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ -1 \\ 6 \end{pmatrix}$$

Is  $b$  in the span of the columns of  $A$ ? In other words, is  $b$  a linear combination of the columns of  $A$ ? Justify your answer.

**Solution.**

Let  $v_1$ ,  $v_2$ , and  $v_3$  be the columns of  $A$ . We are asked to determine whether there are scalars  $x_1$ ,  $x_2$ , and  $x_3$  so that  $x_1v_1 + x_2v_2 + x_3v_3 = b$ , which means

$$\begin{aligned} x_1 + 5x_3 &= 2 \\ -2x_1 + x_2 - 6x_3 &= -1 \\ 2x_2 + 8x_3 &= 6 \end{aligned}$$

We translate the system of linear equations into an augmented matrix, and row reduce it:

$$\left( \begin{array}{ccc|c} 1 & 0 & 5 & 2 \\ -2 & 1 & -6 & -1 \\ 0 & 2 & 8 & 6 \end{array} \right) \xrightarrow{\text{rref}} \left( \begin{array}{ccc|c} 1 & 0 & 5 & 2 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

The right column is not a pivot column, so the system is consistent. Therefore,  $b$  is in the span of the columns of  $A$  (in other words,  $b$  is a linear combination of the columns of  $A$ ).

We weren't asked to solve the equation explicitly, but if we wanted to do so, we would use the RREF of the matrix above to write

$$x_1 = 2 - 5x_3 \quad x_2 = 3 - 4x_3 \quad x_3 = x_3 \quad (x_3 \text{ is free}).$$

In fact, we can take  $x_1 = 2$ ,  $x_2 = 3$ , and  $x_3 = 0$ , to write

$$b = \begin{pmatrix} 2 \\ -1 \\ 6 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} + 0 \begin{pmatrix} 5 \\ -6 \\ 8 \end{pmatrix}.$$

4. Consider the vector equation

$$x \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + y \begin{pmatrix} -2 \\ -1 \\ -1 \end{pmatrix} + z \begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} = \begin{pmatrix} -5 \\ -1 \\ -2 \end{pmatrix}.$$

**Question:** Is there a solution? If so, what is the solution set?

- Formulate this question as an augmented matrix.
- Formulate this question as a system of linear equations.
- What does this mean in terms of spans?
- Answer the question using the [interactive demo](#).
- Answer the question using row reduction.

**Solution.**

a) As an augmented matrix:

$$\left( \begin{array}{ccc|c} 2 & -2 & 3 & -5 \\ 1 & -1 & 0 & -1 \\ 3 & -1 & 4 & -2 \end{array} \right)$$

b) What is the solution set of the following linear system?

$$\begin{aligned} 2x - 2y + 3z &= -5 \\ x - y &= -1 \\ 3x - y + 4z &= -2 \end{aligned}$$

c) There exists a solution if and only if  $\begin{pmatrix} -5 \\ -1 \\ -2 \end{pmatrix}$  is in  $\text{Span} \left\{ \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} -2 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix} \right\}$ .

e) Row reducing yields

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & 3/2 \\ 0 & 1 & 0 & 5/2 \\ 0 & 0 & 1 & -1 \end{array} \right),$$

so  $x = 3/2$ ,  $y = 5/2$ , and  $z = -1$ .

5. Let  $v_1 = \begin{pmatrix} 1 \\ k \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} -1 \\ 4 \end{pmatrix}$ , and  $b = \begin{pmatrix} 1 \\ h \end{pmatrix}$ .

- Find all values of  $h$  and  $k$  so that  $x_1 v_1 + x_2 v_2 = b$  has infinitely many solutions.
- Find all values of  $h$  and  $k$  so that  $b$  is *not* in  $\text{Span}\{v_1, v_2\}$ .
- Find all values of  $h$  and  $k$  so that there is exactly one way to express  $b$  as a linear combination of  $v_1$  and  $v_2$ .

**Solution.**

Each part uses the row-reduction

$$\left( \begin{array}{cc|c} 1 & -1 & 1 \\ k & 4 & h \end{array} \right) \xrightarrow{R_2=R_2-kR_1} \left( \begin{array}{cc|c} 1 & -1 & 1 \\ 0 & 4+k & h-k \end{array} \right).$$

- The system  $(v_1 \ v_2 \mid b)$  has infinitely many solutions if and only if the right column is not a pivot column and there is at least one free variable. This means that  $4+k=0$  and  $h-k=0$ , so  $k=-4$  and  $h=k$ , thus  $\boxed{k=-4 \text{ and } h=-4}$ .
- The right column is a pivot column when  $4+k=0$  and  $h-k \neq 0$ . Thus  $\boxed{k=-4 \text{ and } h \neq -4}$ .
- The system will have a unique solution when the right column is not a pivot column but both other columns are pivot columns. This is when  $4+k \neq 0$ , so  $\boxed{k \neq -4 \text{ and } h \text{ is any real number}}$ .

6. Decide if each of the following statements is true or false. If it is true, prove it; if it is false, provide a counterexample.
- Every set of four or more vectors in  $\mathbf{R}^3$  will span  $\mathbf{R}^3$ .
  - The span of any set contains the zero vector.

**Solution.**

- a) This is **false**. For instance, the vectors

$$\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix} \right\}$$

only span the  $x$ -axis.

- b) This is **true**. We have

$$0 = 0 \cdot v_1 + 0 \cdot v_2 + \cdots + 0 \cdot v_p.$$

Aside: the span of the empty set is equal to  $\{0\}$ , because  $0$  is the empty sum, i.e. the sum with no summands. Indeed, if you add the empty sum to a vector  $v$ , you get  $v +$  (no other summands), which is just  $v$ ; and the only vector which gives you  $v$  when you add it to  $v$ , is  $0$ . (If you find this argument intriguing, you might want to consider taking abstract math courses later on.)

7. Is  $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$  in the span of  $\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$  and  $\begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix}$ ? Justify your answer.

**Solution.**

No. We row-reduce the corresponding augmented matrix to get

$$\left( \begin{array}{cc|c} 0 & 2 & 0 \\ 1 & 3 & 1 \\ 1 & 1 & 0 \end{array} \right) \xrightarrow{RREF} \left( \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

which is inconsistent since it has a pivot in the right column.

8. Acme Widgets, Gizmos, and Doodads has two factories. Factory A makes 10 widgets, 3 gizmos, and 2 doodads every hour, and factory B makes 4 widgets, 1 gizmo, and 1 doodad every hour.
- If factory A runs for  $a$  hours and factory B runs for  $b$  hours, how many widgets, gizmos, and doodads are produced? Express your answer as a vector equation.
  - A customer places an order for 16 widgets, 5 gizmos, and 3 doodads. Can Acme fill the order with no widgets, gizmos, or doodads left over? If so, how many hours do the factories run? If not, why not?

**Solution.**

a) Let  $w$ ,  $g$ , and  $d$  be the number of widgets, gizmos, and doodads produced.

$$\begin{pmatrix} w \\ g \\ d \end{pmatrix} = a \begin{pmatrix} 10 \\ 3 \\ 2 \end{pmatrix} + b \begin{pmatrix} 4 \\ 1 \\ 1 \end{pmatrix}.$$

b) We need to solve the vector equation


$$\begin{pmatrix} 16 \\ 5 \\ 3 \end{pmatrix} = a \begin{pmatrix} 10 \\ 3 \\ 2 \end{pmatrix} + b \begin{pmatrix} 4 \\ 1 \\ 1 \end{pmatrix}.$$

We put it into an augmented matrix and row reduce:

$$\begin{pmatrix} 10 & 4 & | & 16 \\ 3 & 1 & | & 5 \\ 2 & 1 & | & 3 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 3 & 1 & | & 5 \\ 2 & 1 & | & 3 \\ 10 & 4 & | & 16 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 0 & | & 2 \\ 2 & 1 & | & 3 \\ 10 & 4 & | & 16 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 0 & | & 2 \\ 0 & 1 & | & -1 \\ 10 & 4 & | & 16 \end{pmatrix} \\ \rightsquigarrow \begin{pmatrix} 1 & 0 & | & 2 \\ 0 & 1 & | & -1 \\ 0 & 0 & | & 0 \end{pmatrix}$$

These equations are consistent, but they tell us that factory B would have to run for  $-1$  hours! Therefore it can't be done.

9. The diagram below represents traffic in a city.



- a) Write a system of three linear equations whose solution would give the values of  $x_1$ ,  $x_2$ , and  $x_3$ . Do not solve it.
- b) Write the system of equations as a vector equation. Do not solve it.

**Solution.**

a) The number of cars leaving an intersection must equal the number of cars entering.

$$x_3 + 70 = x_1 + 90$$

$$x_1 + x_2 = 160$$

$$x_2 + x_3 = 180.$$

Or:

$$-x_1 + x_3 = 20$$

$$x_1 + x_2 = 160$$

$$x_2 + x_3 = 180.$$

$$\mathbf{b)} \quad x_1 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + x_3 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 20 \\ 160 \\ 180 \end{pmatrix}.$$