

Supplemental problems: §5.1

1. True or false. Answer true if the statement is always true. Otherwise, answer false.
 - a) If A and B are $n \times n$ matrices and A is row equivalent to B , then A and B have the same eigenvalues.
 - b) If A is an $n \times n$ matrix and its eigenvectors form a basis for \mathbf{R}^n , then A is invertible.
 - c) If 0 is an eigenvalue of the $n \times n$ matrix A , then $\text{rank}(A) < n$.
 - d) The diagonal entries of an $n \times n$ matrix A are its eigenvalues.
 - e) If A is invertible and 2 is an eigenvalue of A , then $\frac{1}{2}$ is an eigenvalue of A^{-1} .
 - f) If $\det(A) = 0$, then 0 is an eigenvalue of A .

2. In this problem, you need not explain your answers; just circle the correct one(s).

Let A be an $n \times n$ matrix.

- a) Which **one** of the following statements is correct?

1. An eigenvector of A is a vector v such that $Av = \lambda v$ for a nonzero scalar λ .
2. An eigenvector of A is a nonzero vector v such that $Av = \lambda v$ for a scalar λ .
3. An eigenvector of A is a nonzero scalar λ such that $Av = \lambda v$ for some vector v .
4. An eigenvector of A is a nonzero vector v such that $Av = \lambda v$ for a nonzero scalar λ .

- b) Which **one** of the following statements is **not** correct?

1. An eigenvalue of A is a scalar λ such that $A - \lambda I$ is not invertible.
2. An eigenvalue of A is a scalar λ such that $(A - \lambda I)v = 0$ has a solution.
3. An eigenvalue of A is a scalar λ such that $Av = \lambda v$ for a nonzero vector v .
4. An eigenvalue of A is a scalar λ such that $\det(A - \lambda I) = 0$.

3. Find a basis \mathcal{B} for the (-1) -eigenspace of $Z = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}$

4. Suppose A is an $n \times n$ matrix satisfying $A^2 = 0$. Find all eigenvalues of A . Justify your answer.

5. Match the statements (i)-(v) with the corresponding statements (a)-(e). All matrices are 3×3 . There is a unique correspondence. Justify the correspondences in words.

(i) $Ax = \begin{pmatrix} 5 \\ 1 \\ 2 \end{pmatrix}$ has a unique solution.

(ii) The transformation $T(v) = Av$ fixes a nonzero vector.

(iii) A is obtained from B by subtracting the third row of B from the first row of B .

(iv) The columns of A and B are the same; except that the first, second and third columns of A are respectively the first, third, and second columns of B .

(v) The columns of A , when added, give the zero vector.

(a) 0 is an eigenvalue of A .

(b) A is invertible.

(c) $\det(A) = \det(B)$

(d) $\det(A) = -\det(B)$

(e) 1 is an eigenvalue of A .

Supplemental problems: §5.2

1. True or false. If the statement is always true, answer true and justify why it is true. Otherwise, answer false and give an example that shows it is false.
 - a) If A and B are $n \times n$ matrices with the same eigenvectors, then A and B have the same characteristic polynomial.
 - b) If A is a 3×3 matrix with characteristic polynomial $-\lambda^3 + \lambda^2 + \lambda$, then A is invertible.
2. Find all values of a so that $\lambda = 1$ an eigenvalue of the matrix A below.

$$A = \begin{pmatrix} 3 & -1 & 0 & a \\ a & 2 & 0 & 4 \\ 2 & 0 & 1 & -2 \\ 13 & a & -2 & -7 \end{pmatrix}$$