

MATH 1553, FALL 2019
SAMPLE MIDTERM 3B: COVERS 4.1 THROUGH 5.5

| | | | |
|-------------|--|----------------|--|
| Name | | Section | |
|-------------|--|----------------|--|

Please **read all instructions** carefully before beginning.

- Write your name on the front of each page (not just the cover page!).
- The maximum score on this exam is 50 points, and you have 50 minutes to complete this exam.
- There are no calculators or aids of any kind (notes, text, etc.) allowed.
- As always, RREF means “reduced row echelon form”.
- Show your work, unless instructed otherwise. A correct answer without appropriate work will receive little or no credit! If you cannot fit your work on the front side of the page, use the back side of the page and indicate that you are using the back side.
- We will hand out loose scrap paper, but it **will not be graded** under any circumstances. All work must be written on the exam itself.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 50 minutes, without notes or distractions.

The exam is not designed to test material from the previous midterm on its own. However, knowledge of the material prior to section §4.1 is necessary for everything we do for the rest of the semester, so it is fair game for the exam as it applies to §§4.1 through 5.5.

This page was intentionally left blank.

Problem 1.

Answer true if the statement is *always* true. Otherwise, answer false. In every case, assume that the entries of the matrix A are real numbers.


- a) **T** **F** If A is the 3×3 matrix satisfying $Ae_1 = e_2$, $Ae_2 = e_3$, and $Ae_3 = e_1$, then $\det(A) = 1$.
- b) **T** **F** If A is an $n \times n$ matrix and $\det(A) = 2$, then 2 is an eigenvalue of A .
- c) **T** **F** If A and B are $n \times n$ matrices with $\det(A) = 0$ and $\det(B) = 0$, then $\det(A + B) = 0$.
- d) **T** **F** If A is an $n \times n$ matrix and v and w are eigenvectors of A , then $v + w$ is also an eigenvector of A .
- e) **T** **F** It is possible for a lower-triangular matrix A to have a non-real complex eigenvalue.

Extra space for scratch work on problem 1


Problem 2.

Short answer. Show your work on part (c). In every case, the entries of each matrix must be real numbers.

- a) Write a 2×2 matrix A which is invertible but not diagonalizable.
- b) Write a 2×2 matrix A for which $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are eigenvectors corresponding to the same eigenvalue.
- c) Find the area of the parallelogram drawn below (the grid marks are spaced one unit apart).



- d) Write a 3×3 matrix A with only one real eigenvalue $\lambda = 4$, such that the 4-eigenspace for A is a two-dimensional plane in \mathbb{R}^3 .
- e) Suppose that $A = C \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix} C^{-1}$, where C has columns v_1 and v_2 . Given x and y in the picture below, draw the vectors Ax and Ay .



Extra space for work on problem 2

Problem 3.

Parts (a) and (b) are unrelated.

a) Consider the matrix

$$A = \begin{pmatrix} 3 & -7 \\ 1 & -1 \end{pmatrix}$$

Find all eigenvalues of A . Simplify your answer. For the eigenvalue with negative imaginary part, find an eigenvector.

b) Let $A = \begin{pmatrix} 7 & -8 \\ 4 & -5 \end{pmatrix}$. Find a formula for A^n and simplify your answer completely.

Extra space for work on problem 3


Extra space for work on problem 4

Problem 5.

Parts (a) and (b) are not related.

a) Find $\det(A^3)$ if $A = \begin{pmatrix} 1 & -3 & 4 & 2 \\ 0 & 0 & -2 & 0 \\ 0 & 1 & 2 & 3 \\ 2 & 0 & -1 & 20 \end{pmatrix}$.

b) Find the 2×2 matrix A whose eigenspaces are drawn below. Fully simplify your answer. (to be clear: the dashed line is the (-2) -eigenspace).



Extra space for work on problem 5