

MATH 1553, FALL 2019
SAMPLE MIDTERM 3A: COVERS 4.1 THROUGH 5.5

Name		Section	
-------------	--	----------------	--

Please **read all instructions** carefully before beginning.

- Write your name on the front of each page (not just the cover page!).
- The maximum score on this exam is 50 points, and you have 50 minutes to complete this exam.
- There are no calculators or aids of any kind (notes, text, etc.) allowed.
- As always, RREF means “reduced row echelon form”.
- Show your work, unless instructed otherwise. A correct answer without appropriate work will receive little or no credit! If you cannot fit your work on the front side of the page, use the back side of the page and indicate that you are using the back side.
- We will hand out loose scrap paper, but it **will not be graded** under any circumstances. All work must be written on the exam itself.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 50 minutes, without notes or distractions.

The exam is not designed to test material from the previous midterm on its own. However, knowledge of the material prior to section §4.1 is necessary for everything we do for the rest of the semester, so it is fair game for the exam as it applies to §§4.1 through 5.5.

This page was intentionally left blank.

Problem 1.

True or false. Circle **T** if the statement is *always* true.

Otherwise, circle **F**. You do not need to show work or justify your answer.

- a) **T** **F** If A is an $n \times n$ matrix, then the determinant of A is the same as the determinant of the RREF of A .
- b) **T** **F** If A is a 3×3 matrix with characteristic polynomial
$$\det(A - \lambda I) = (1 - \lambda)(-1 - \lambda)^2,$$
then A must be invertible.
- c) **T** **F** Suppose A is an $n \times n$ matrix and λ is an eigenvalue of A . If v and w are two different eigenvectors of A corresponding to the eigenvalue λ , then $v - w$ is an eigenvector of A .
- d) **T** **F** If A and B are 3×3 matrices that have the same eigenvalues and the same algebraic multiplicity for each eigenvalue, then $A = B$.
- e) **T** **F** If A is a 4×4 matrix, then $\det(-A) = \det(A)$.

Extra space for scratch work on problem 1

Problem 2.

You do not need to show your work or justify your answers. Each part is worth 2 points except (e), which is worth 4 points.


a) Complete the following definition (be mathematically precise!):

Suppose A is an $n \times n$ matrix and λ is a real number. We say λ is an *eigenvalue* of A if...

b) Suppose $\det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} = 3$. Find $\det \begin{pmatrix} -4a + d & -4b + e & -4c + f \\ a & b & c \\ g & h & i \end{pmatrix}$.

c) Write a 2×2 matrix which is neither diagonalizable nor invertible.

d) Let A be the matrix which implements reflection in \mathbf{R}^2 across the line $y = -x/2$. In the graph below, clearly draw one eigenvector in each eigenspace of A (you don't need to write the eigenvalues of A)



e) Suppose A is an $n \times n$ matrix and $\det(A) = 0$.

Which of the following statements must be true? Circle all that apply.

(i) $\dim(\text{Nul}(A)) \geq 1$.

(ii) The equation $Ax = 0$ has only the trivial solution $x = 0$.

(iii) $\lambda = 0$ is an eigenvalue of A .

(iv) The equation $Ax = b$ must be inconsistent for at least one b in \mathbf{R}^n .

Extra space for work on problem 2

Problem 3.

Parts (a) and (b) are unrelated.

- a) Let $A = \begin{pmatrix} 5 & 5 \\ -2 & -1 \end{pmatrix}$. Find the complex eigenvalues of A . For the eigenvalue with positive imaginary part, find one corresponding eigenvector.

- b) Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 0 & 3 \\ 2 & c & c & 1 \\ 3 & 0 & 0 & 4 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

Find all values of c so that $\det(A) = 4$.

Extra space for work on problem 3

Problem 4.

$$\text{Let } A = \begin{pmatrix} 3 & 0 & -2 \\ 2 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}.$$

a) Find all eigenvalues of A .

b) Find a basis for each eigenspace of A .

c) Is A diagonalizable? If so, write an invertible 3×3 matrix C and a diagonal matrix D so that $A = CDC^{-1}$. If not, justify why A is not diagonalizable.

Extra space for work on problem 4

Problem 5.

Parts (a) and (b) are unrelated. No work is required on (a).

a) Suppose $A = \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} -1/3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix}^{-1}$. Which of the following are true? Circle all that apply.

(i) Every nonzero vector in \mathbf{R}^2 is an eigenvector of A .

(ii) Repeated multiplication by A pushes vectors toward $\text{Span} \left\{ \begin{pmatrix} 2 \\ 3 \end{pmatrix} \right\}$.

(iii) If $x = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$, then $A^n x$ approaches the zero vector as n becomes very large.

(iv) The eigenvalues of A are $-\frac{1}{3}$ and 1.

b) Suppose A is a 2×2 matrix satisfying $\text{Tr}(A) = 6$ and $\det(A) = 9$. Must A be invertible? Must A be diagonalizable? Justify your answers.

Extra space for work on problem 5