

Section 7.2

Orthogonal Complements

Orthogonal Complements

Definition

Let W be a subspace of \mathbf{R}^n . Its **orthogonal complement** is

$$W^\perp = \{v \text{ in } \mathbf{R}^n \mid v \cdot w = 0 \text{ for all } w \text{ in } W\} \quad \text{read "W perp".}$$




W^\perp is orthogonal complement
 A^T is transpose

Pictures:

The orthogonal complement of a **line** in \mathbf{R}^2 is the perpendicular **line**. [interactive]

The orthogonal complement of a **line** in \mathbf{R}^3 is the perpendicular **plane**. [interactive]

The orthogonal complement of a **plane** in \mathbf{R}^3 is the perpendicular **line**. [interactive]



Poll

Let W be a 2-plane in \mathbf{R}^4 . How would you describe W^\perp ?

- A. The zero space $\{0\}$.
- B. A line in \mathbf{R}^4 .
- C. A plane in \mathbf{R}^4 .
- D. A 3-dimensional space in \mathbf{R}^4 .
- E. All of \mathbf{R}^4 .

For example, if W is the xy -plane, then W^\perp is the zw -plane:

$$\begin{pmatrix} x \\ y \\ 0 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 0 \\ z \\ w \end{pmatrix} = 0.$$

Orthogonal Complements

Basic properties

Let W be a subspace of \mathbf{R}^n .

Facts:

1. W^\perp is also a subspace of \mathbf{R}^n
2. $(W^\perp)^\perp = W$
3. $\dim W + \dim W^\perp = n$
4. If $W = \text{Span}\{v_1, v_2, \dots, v_m\}$, then

$$\begin{aligned}W^\perp &= \text{all vectors orthogonal to each } v_1, v_2, \dots, v_m \\ &= \{x \text{ in } \mathbf{R}^n \mid x \cdot v_i = 0 \text{ for all } i = 1, 2, \dots, m\} \\ &= \text{Nul} \begin{pmatrix} -v_1^T & - \\ -v_2^T & - \\ \vdots & \\ -v_m^T & - \end{pmatrix}.\end{aligned}$$

Let's check 1.

- ▶ Is 0 in W^\perp ? Yes: $0 \cdot w = 0$ for any w in W .
- ▶ Suppose x, y are in W^\perp . So $x \cdot w = 0$ and $y \cdot w = 0$ for all w in W . Then $(x + y) \cdot w = x \cdot w + y \cdot w = 0 + 0 = 0$ for all w in W . So $x + y$ is also in W^\perp .
- ▶ Suppose x is in W^\perp . So $x \cdot w = 0$ for all w in W . If c is a scalar, then $(cx) \cdot w = c(x \cdot w) = c(0) = 0$ for any w in W . So cx is in W^\perp .

Orthogonal Complements

Computation

Problem: if $W = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$, compute W^\perp .

By property 4, we have to find the null space of the matrix whose rows are $(1 \ 1 \ -1)$ and $(1 \ 1 \ 1)$, which we did before:

$$\text{Nul} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \end{pmatrix} = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

[interactive]

$$\text{Span}\{v_1, v_2, \dots, v_m\}^\perp = \text{Nul} \begin{pmatrix} -v_1^T & - \\ -v_2^T & - \\ \vdots & \\ -v_m^T & - \end{pmatrix}$$

Orthogonal Complements

Row space, column space, null space

Definition

The **row space** of an $m \times n$ matrix A is the span of the *rows* of A . It is denoted $\text{Row } A$. Equivalently, it is the column space of A^T :

$$\text{Row } A = \text{Col } A^T.$$

It is a subspace of \mathbf{R}^n .

We showed before that if A has rows $v_1^T, v_2^T, \dots, v_m^T$, then

$$\text{Span}\{v_1, v_2, \dots, v_m\}^\perp = \text{Nul } A.$$

Hence we have shown:

Fact: $(\text{Row } A)^\perp = \text{Nul } A$.

Replacing A by A^T , and remembering $\text{Row } A^T = \text{Col } A$:

Fact: $(\text{Col } A)^\perp = \text{Nul } A^T$.

Using property 2 and taking the orthogonal complements of both sides, we get:

Fact: $(\text{Nul } A)^\perp = \text{Row } A$ and $\text{Col } A = (\text{Nul } A^T)^\perp$.

Orthogonal Complements of Most of the Subspaces We've Seen

For any vectors v_1, v_2, \dots, v_m :

$$\text{Span}\{v_1, v_2, \dots, v_m\}^\perp = \text{Nul} \begin{pmatrix} -v_1^T & - \\ -v_2^T & - \\ \vdots & \\ -v_m^T & - \end{pmatrix}$$

For any matrix A :

$$\text{Row } A = \text{Col } A^T$$

and

$$(\text{Row } A)^\perp = \text{Nul } A \quad \text{Row } A = (\text{Nul } A)^\perp$$

$$(\text{Col } A)^\perp = \text{Nul } A^T \quad \text{Col } A = (\text{Nul } A^T)^\perp$$


For any other subspace W , first find a basis v_1, \dots, v_m , then use the above trick to compute $W^\perp = \text{Span}\{v_1, \dots, v_m\}^\perp$.

Section 7.3

Orthogonal Projections

Best Approximation

Suppose you measure a data point x which you know for theoretical reasons must lie on a subspace W .



Due to measurement error, though, the measured x is not actually in W . Best approximation: y is the *closest* point to x on W .

How do you know that y is the closest point? The vector from y to x is orthogonal to W : it is in the *orthogonal complement* W^\perp .

Orthogonal Decomposition

Theorem

Every vector x in \mathbf{R}^n can be written as

$$x = x_W + x_{W^\perp}$$

for unique vectors x_W in W and x_{W^\perp} in W^\perp .


The equation $x = x_W + x_{W^\perp}$ is called the **orthogonal decomposition** of x (with respect to W).

The vector x_W is the **orthogonal projection** of x onto W .

The vector x_W is the *closest vector to x on W* .

[interactive 1]

[interactive 2]



Orthogonal Decomposition

Justification

Theorem

Every vector x in \mathbf{R}^n can be written as

$$x = x_W + x_{W^\perp}$$

for unique vectors x_W in W and x_{W^\perp} in W^\perp .

Why?

Uniqueness: suppose $x = x_W + x_{W^\perp} = x'_W + x'_{W^\perp}$ for x_W, x'_W in W and $x_{W^\perp}, x'_{W^\perp}$ in W^\perp . Rewrite:

$$x_W - x'_W = x'_{W^\perp} - x_{W^\perp}.$$

The left side is in W , and the right side is in W^\perp , so they are both in $W \cap W^\perp$. But the only vector that is perpendicular to itself is the zero vector! Hence

$$\begin{aligned} 0 &= x_W - x'_W \implies x_W = x'_W \\ 0 &= x_{W^\perp} - x'_{W^\perp} \implies x_{W^\perp} = x'_{W^\perp} \end{aligned}$$

Existence: We will compute the orthogonal decomposition later using orthogonal projections.

Orthogonal Decomposition


Example

Let W be the xy -plane in \mathbf{R}^3 . Then W^\perp is the z -axis.

$$x = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} \implies x_W = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} \quad x_{W^\perp} = \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix}.$$

$$x = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \implies x_W = \begin{pmatrix} a \\ b \\ 0 \end{pmatrix} \quad x_{W^\perp} = \begin{pmatrix} 0 \\ 0 \\ c \end{pmatrix}.$$

This is just decomposing a vector into a “horizontal” component (in the xy -plane) and a “vertical” component (on the z -axis).



[interactive]

Orthogonal Decomposition

Computation?

Problem: Given x and W , how do you compute the decomposition $x = x_W + x_{W^\perp}$?

Observation: It is enough to compute x_W , because $x_{W^\perp} = x - x_W$.

Theorem (The $A^T A$ Trick)

Let W be a subspace of \mathbf{R}^n , let v_1, v_2, \dots, v_m be a spanning set for W (e.g., a basis), and let

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_m \\ | & | & \cdots & | \end{pmatrix}.$$

Then for any x in \mathbf{R}^n , the matrix equation

$$A^T A v = A^T x \quad (\text{in the unknown vector } v)$$

is consistent, and $x_W = Av$ for any solution v .

Recipe for Computing $x = x_W + x_{W^\perp}$

- ▶ Write W as a column space of a matrix A .
- ▶ Find a solution v of $A^T A v = A^T x$ (by row reducing).
- ▶ Then $x_W = Av$ and $x_{W^\perp} = x - x_W$.

The $A^T A$ Trick

Example

Problem: Compute the orthogonal projection of a vector $x = (x_1, x_2, x_3)$ in \mathbf{R}^3 onto the xy -plane.

First we need a basis for the xy -plane: let's choose

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \rightsquigarrow \quad A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Then

$$A^T A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I_2 \quad A^T \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

Then $A^T A v = v$ and $A^T x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$, so the only solution of $A^T A v = A^T x$ is $v = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Therefore,

$$x_W = A v = A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix}.$$

The $A^T A$ Trick

Another Example

Problem: Let

$$x = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad W = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \text{ in } \mathbf{R}^3 \mid x_1 - x_2 + x_3 = 0 \right\}.$$

Compute the distance from x to W .

The distance from x to W is $\|x_{W^\perp}\|$, so we need to compute the orthogonal projection. First we need a basis for $W = \text{Nul} \begin{pmatrix} 1 & -1 & 1 \end{pmatrix}$. This matrix is in RREF, so the parametric form of the solution set is

$$\begin{array}{l} x_1 = x_2 - x_3 \\ x_2 = x_2 \\ x_3 = x_3 \end{array} \quad \begin{array}{l} \text{PVF} \\ \rightsquigarrow \end{array} \quad \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}.$$

Hence we can take a basis to be

$$\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\} \rightsquigarrow A = \begin{pmatrix} 1 & -1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$$

The $A^T A$ Trick

Another Example, Continued

Problem: Let

$$x = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad W = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \text{ in } \mathbf{R}^3 \mid x_1 - x_2 + x_3 = 0 \right\}.$$

Compute the distance from x to W .

We compute

$$A^T A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \quad A^T x = \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

To solve $A^T A v = A^T x$ we form an augmented matrix and row reduce:

$$\left(\begin{array}{cc|c} 2 & -1 & 3 \\ -1 & 2 & 2 \end{array} \right) \xrightarrow{\text{RREF}} \left(\begin{array}{ccc} 1 & 0 & 8/3 \\ 0 & 1 & 7/3 \end{array} \right) \rightsquigarrow v = \frac{1}{3} \begin{pmatrix} 8 \\ 7 \end{pmatrix}.$$

$$x_W = Av = \frac{1}{3} \begin{pmatrix} 1 \\ 8 \\ 7 \end{pmatrix} \quad x_{W^\perp} = x - x_W = \frac{1}{3} \begin{pmatrix} 2 \\ -2 \\ 2 \end{pmatrix}.$$

The distance is $\|x_{W^\perp}\| = \frac{1}{3}\sqrt{4+4+4} \approx 1.155$.

[interactive]

The $A^T A$ trick

Proof

Theorem (The $A^T A$ Trick)

Let W be a subspace of \mathbf{R}^n , let v_1, v_2, \dots, v_m be a spanning set for W (e.g., a basis), and let

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_m \\ | & | & \cdots & | \end{pmatrix}.$$

Then for any x in \mathbf{R}^n , the matrix equation

$$A^T A v = A^T x \quad (\text{in the unknown vector } v)$$

is consistent, and $x_W = A v$ for any solution v .

Proof: Let $x = x_W + x_{W^\perp}$. Then x_{W^\perp} is in $W^\perp = \text{Nul}(A^T)$, so $A^T x_{W^\perp} = 0$. Hence

$$A^T x = A^T (x_W + x_{W^\perp}) = A^T x_W + A^T x_{W^\perp} = A^T x_W.$$

Since x_W is in $W = \text{Span}\{v_1, v_2, \dots, v_m\}$, we can write

$$x_W = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

If $v = (c_1, c_2, \dots, c_m)$ then $A v = x_W$, so

$$A^T x = A^T x_W = A^T A v.$$

Orthogonal Projection onto a Line

Problem: Let $L = \text{Span}\{u\}$ be a line in \mathbf{R}^n and let x be a vector in \mathbf{R}^n . Compute x_L .


We have to solve $u^T uv = u^T x$, where u is an $n \times 1$ matrix. But $u^T u = u \cdot u$ and $u^T x = u \cdot x$ are scalars, so

$$v = \frac{u \cdot x}{u \cdot u} \implies x_L = uv = \frac{u \cdot x}{u \cdot u} u.$$

Projection onto a Line

The projection of x onto a line $L = \text{Span}\{u\}$ is

$$x_L = \frac{u \cdot x}{u \cdot u} u \quad x_{L^\perp} = x - x_L.$$



Orthogonal Projection onto a Line


Example

Problem: Compute the orthogonal projection of $x = \begin{pmatrix} -6 \\ 4 \end{pmatrix}$ onto the line L spanned by $u = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$, and find the distance from x to L .

$$x_L = \frac{x \cdot u}{u \cdot u} u = \frac{-18 + 8}{9 + 4} \begin{pmatrix} 3 \\ 2 \end{pmatrix} = -\frac{10}{13} \begin{pmatrix} 3 \\ 2 \end{pmatrix} \quad x_{L^\perp} = x - x_L = \frac{1}{13} \begin{pmatrix} -48 \\ 72 \end{pmatrix}.$$

The distance from x to L is

$$\|x_{L^\perp}\| = \frac{1}{13} \sqrt{48^2 + 72^2} \approx 6.656.$$



[interactive]

Let W be a subspace of \mathbf{R}^n .

- ▶ The **orthogonal complement** W^\perp is the set of all vectors orthogonal to everything in W .
- ▶ We have $(W^\perp)^\perp = W$ and $\dim W + \dim W^\perp = n$.
- ▶ $\text{Row } A = \text{Col } A^T$, $(\text{Row } A)^\perp = \text{Nul } A$, $\text{Row } A = (\text{Nul } A)^\perp$,
 $(\text{Col } A)^\perp = \text{Nul } A^T$, $\text{Col } A = (\text{Nul } A^T)^\perp$.
- ▶ **Orthogonal decomposition:** any vector x in \mathbf{R}^n can be written in a unique way as $x = x_W + x_{W^\perp}$ for x_W in W and x_{W^\perp} in W^\perp . The vector x_W is the **orthogonal projection** of x onto W .
- ▶ The vector x_W is the *closest point to x in W* : it is the *best approximation*.
- ▶ The *distance* from x to W is $\|x_{W^\perp}\|$.
- ▶ If $W = \text{Col } A$ then to compute x_W , solve the equation $A^T A v = A^T x$; then $x_W = A v$.
- ▶ If $W = L = \text{Span}\{u\}$ is a line then $x_L = \frac{u \cdot x}{u \cdot u} u$.