

## Math 1553 Supplement, Chapter 7

1. True or false. If the statement is always true, answer true. Otherwise, answer false. Justify your answer.
- a) Suppose  $W = \text{Span}\{w\}$  for some vector  $w \neq 0$ , and suppose  $v$  is a vector orthogonal to  $w$ . Then the orthogonal projection of  $v$  onto  $W$  is the zero vector.
  - b) Suppose  $W$  is a subspace of  $\mathbf{R}^n$  and  $x$  is a vector in  $\mathbf{R}^n$ . If  $x$  is not in  $W$ , then  $x - x_W$  is not zero.
  - c) Suppose  $W$  is a subspace of  $\mathbf{R}^n$  and  $x$  is in both  $W$  and  $W^\perp$ . Then  $x = 0$ .
  - d) Suppose  $\hat{x}$  is a least squares solution to  $Ax = b$ . Then  $\hat{x}$  is the closest vector to  $b$  in the column space of  $A$ .

### Solution.

- a) True. Since  $v \in W^\perp$ , its projection onto  $W$  is zero.
  - b) True. If  $x$  is not in  $W$  then  $x \neq x_W$ , so  $x - x_W$  is not zero.
  - c) True. Since  $x$  is in  $W$  and  $W^\perp$  it is orthogonal to itself, so  $\|x\|^2 = x \cdot x = 0$ . The length of  $x$  is zero, which means every entry of  $x$  is zero, hence  $x = 0$ .
  - d) False:  $A\hat{x}$  is the closest vector to  $b$  in  $\text{Col } A$ .
2. Let  $W = \text{Span}\{v_1, v_2\}$ , where  $v_1 = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}$  and  $v_2 = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ .

- a) Find the closest point  $w$  in  $W$  to  $x = \begin{pmatrix} 0 \\ 14 \\ -4 \end{pmatrix}$ .

Let  $A = \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix}$ . We solve  $A^T A v = A^T x$ .

$$A^T A = \begin{pmatrix} 6 & 6 \\ 6 & 14 \end{pmatrix} \quad A^T \begin{pmatrix} 0 \\ 14 \\ -4 \end{pmatrix} = \begin{pmatrix} 24 \\ 16 \end{pmatrix}.$$

We find  $\left( \begin{array}{cc|c} 6 & 6 & 24 \\ 6 & 14 & 16 \end{array} \right) \xrightarrow{RREF} \left( \begin{array}{cc|c} 1 & 0 & 5 \\ 0 & 1 & -1 \end{array} \right)$ , so  $v = \begin{pmatrix} 5 \\ -1 \end{pmatrix}$  and therefore

$$w = Av = \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 5 \\ -1 \end{pmatrix} = \begin{pmatrix} -6 \\ 8 \\ 2 \end{pmatrix}.$$

b) Find the distance from  $w$  to  $\begin{pmatrix} 0 \\ 14 \\ -4 \end{pmatrix}$ .

$$\|x - w\| = \left\| \begin{pmatrix} 0 \\ 14 \\ -4 \end{pmatrix} - \begin{pmatrix} -6 \\ 8 \\ 2 \end{pmatrix} \right\| = \left\| \begin{pmatrix} 6 \\ 6 \\ -6 \end{pmatrix} \right\| = \sqrt{36 + 36 + 36} = \sqrt{108} = 6\sqrt{3}.$$

c) Find the standard matrix for the orthogonal projection onto  $\text{Span}\{v_1\}$ .

$$B = \frac{1}{v_1 \cdot v_1} v_1 v_1^T = \frac{1}{(-1)^2 + 2^2 + 1^2} \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} \begin{pmatrix} -1 & 2 & 1 \end{pmatrix} = \frac{1}{6} \begin{pmatrix} 1 & -2 & -1 \\ -2 & 4 & 2 \\ -1 & 2 & 1 \end{pmatrix}$$

d) Find the standard matrix for the orthogonal projection onto  $W$ .

Let  $A = \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix}$ . Since the columns of  $A$  are linearly independent, our projection matrix is  $A(A^T A)^{-1} A^T$ . We already computed  $A^T A$  in part (a), so our matrix is

$$\begin{aligned} A(A^T A)^{-1} A^T &= \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 6 & 6 \\ 6 & 14 \end{pmatrix}^{-1} \begin{pmatrix} -1 & 2 & 1 \\ 1 & 2 & 3 \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 6 & 6 \\ 6 & 14 \end{pmatrix}^{-1} \begin{pmatrix} -1 & 2 & 1 \\ 1 & 2 & 3 \end{pmatrix} \\ &= \frac{1}{48} \begin{pmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 14 & -6 \\ -6 & 6 \end{pmatrix} \begin{pmatrix} -1 & 2 & 1 \\ 1 & 2 & 3 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix}. \end{aligned}$$