

MATH 1553, SPRING 2019
SAMPLE MIDTERM 1: THROUGH SECTION 3.5

Name	
-------------	--

Please **read all instructions** carefully before beginning.

- You have 50 minutes to complete this exam.
- There are no aids of any kind (calculators, notes, text, etc.) allowed.
- Please show your work unless specified otherwise. A correct answer without appropriate work may be given little or no credit.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 50 minutes, without notes or distractions.

Problem 1.

[Parts a) through e) are worth 2 points each]

a) Compute: $\begin{pmatrix} 3 & 2 \\ -2 & 0 \\ 1 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ -3 \end{pmatrix} =$

The remaining problems are True or false. Circle **T** if the statement is **always** true, and circle **F** otherwise. You do not need to justify your answer.

b) **T** **F** The matrix $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ is in reduced row echelon form.

c) **T** **F** If the augmented matrix corresponding to a linear system of equations has a pivot in every row, then the system is consistent.

d) **T** **F** If A is an $m \times n$ matrix and $Ax = 0$ has a unique solution, then $Ax = b$ is consistent for every b in \mathbf{R}^m .

e) **T** **F** The equation $x_1 - \sqrt{5}x_2 = 10 - \pi^2x_3$ is a linear equation in x_1, x_2, x_3 .

Problem 2.

Parts (a) and (b) are 2 points each. Parts (c) and (d) are 3 points each.

a) If A is a 2×3 matrix with 2 pivots, then the set of solutions to $Ax = 0$ is a:


(circle one answer) **point** **line** **plane** **3-plane**

in:

(circle one answer) **R** **\mathbf{R}^2** **\mathbf{R}^3** .

b) Write a vector equation which represents an inconsistent system of two linear equations in x_1 and x_2 .

c) For some 2×2 matrix A and vector b in \mathbf{R}^2 , the solution set of $Ax = b$ is drawn below. Draw the solution set of $Ax = 0$.



d) If b, v, w are vectors in \mathbf{R}^3 and $\text{Span}\{b, v, w\} = \mathbf{R}^3$, is it possible that b is in $\text{Span}\{v, w\}$? Justify your answer.

Problem 3.

[12 points]

- a) Johnny Rico believes that the secret to the universe can be found in the system of two linear equations in x and y given by

$$\begin{aligned}x - y &= h \\ 3x + hy &= -9\end{aligned}$$

where h is a real number. Find all values of h (if any) which make the system have infinitely many solutions. If there is no such h , justify why.

- b) Find all values of k (if any) so that the vectors below are not linearly independent.

$$\begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 6 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ k \\ 1 \end{pmatrix}.$$

Problem 4.

[11 points]

- a) Find the parametric form of the general solution of the following system of equations. Clearly indicate which variables (if any) are free variables.

$$\begin{aligned}x_1 + 2x_2 + 2x_3 - x_4 &= 4 \\2x_1 + 4x_2 + x_3 - 2x_4 &= -1 \\-x_1 - 2x_2 - x_3 + x_4 &= -1\end{aligned}$$

- b) Write the set of solutions to

$$\begin{aligned}x_1 + 2x_2 + 2x_3 - x_4 &= 0 \\2x_1 + 4x_2 + x_3 - 2x_4 &= 0 \\-x_1 - 2x_2 - x_3 + x_4 &= 0\end{aligned}$$

in parametric vector form.

Problem 5.

[7 points]

Write an augmented matrix corresponding to a system of two linear equations in three variables x_1, x_2, x_3 , whose solution set is the span of $\begin{pmatrix} -4 \\ 1 \\ 0 \end{pmatrix}$.

Briefly justify your answer.

[Scratch work]