

## Math 1553 Worksheet §2.2, §2.3

### Solutions

1. Is it possible for a linear system to have a unique solution if it has more equations than variables? If yes, give an example. If no, justify why it is impossible.

#### Solution.

It is possible. One example is the system below, which has unique solution  $x = 5$ ,  $y = 2$ :

$$\begin{aligned}x + y &= 7 \\x - y &= 3 \\2x + 2y &= 14.\end{aligned}$$

2. a) Which of the following matrices are in row echelon form? Which are in reduced row echelon form?
- b) For the matrices in row echelon form, which entries are the pivots? What are the pivot columns?

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1 & 1 \\ 0 & 2 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 4 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

#### Solution.

The first, second, and fourth matrices are in reduced row echelon form; the third matrix is in row echelon form. The pivots are in red; the other entries in the pivot columns are in blue.

3. Find the parametric form of the solutions of following system of equations in  $x_1$ ,  $x_2$ , and  $x_3$  by putting an augmented matrix into reduced row echelon form. State which variables (if any) are free variables. Describe the solution set geometrically.

$$\begin{aligned}x_1 + 3x_2 + x_3 &= 1 \\ -4x_1 - 9x_2 + 2x_3 &= -1 \\ -3x_2 - 6x_3 &= -3.\end{aligned}$$

**Solution.**

$$\begin{aligned}\left(\begin{array}{ccc|c} 1 & 3 & 1 & 1 \\ -4 & -9 & 2 & -1 \\ 0 & -3 & -6 & -3 \end{array}\right) &\xrightarrow{R_2=R_2+4R_1} \left(\begin{array}{ccc|c} 1 & 3 & 1 & 1 \\ 0 & 3 & 6 & 3 \\ 0 & -3 & -6 & -3 \end{array}\right) \\ &\xrightarrow{R_3=R_3+R_2} \left(\begin{array}{ccc|c} 1 & 3 & 1 & 1 \\ 0 & 3 & 6 & 3 \\ 0 & 0 & 0 & 0 \end{array}\right) \\ &\xrightarrow{R_1=R_1-R_2} \left(\begin{array}{ccc|c} 1 & 0 & -5 & -2 \\ 0 & 3 & 6 & 3 \\ 0 & 0 & 0 & 0 \end{array}\right) \\ &\xrightarrow{R_2=R_2\div 3} \left(\begin{array}{ccc|c} 1 & 0 & -5 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 \end{array}\right).\end{aligned}$$

The variables  $x_1$  and  $x_2$  correspond to pivot columns, but  $x_3$  is free.

$$x_1 = -2 + 5x_3, \quad x_2 = 1 - 2x_3, \quad x_3 = x_3 \quad (x_3 \text{ real}).$$

This consistent system in three variables has one free variable, so the solution set is a line in  $\mathbf{R}^3$ .