

# Chapter 6

## Orthogonality

# Section 6.1

Dot Products and Orthogonality


## Orientation

**Recall:** This course is about learning to:

- ▶ Solve the matrix equation  $Ax = b$
- ▶ Solve the matrix equation  $Ax = \lambda x$
- ▶ Almost solve the equation  $Ax = b$

We are now aiming at the last topic.

**Idea:** In the real world, data is imperfect. Suppose you measure a data point  $x$  which you know for theoretical reasons must lie on a plane spanned by two vectors  $u$  and  $v$ .



Due to measurement error, though, the measured  $x$  is not actually in  $\text{Span}\{u, v\}$ . In other words, the equation  $au + bv = x$  has no solution. What do you do? The real value is probably the *closest* point to  $x$  on  $\text{Span}\{u, v\}$ . Which point is that?

## The Dot Product

We need a notion of *angle* between two vectors, and in particular, a notion of *orthogonality* (i.e. when two vectors are perpendicular). This is the purpose of the dot product.

### Definition

The **dot product** of two vectors  $x, y$  in  $\mathbf{R}^n$  is

$$x \cdot y = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \cdot \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} \stackrel{\text{def}}{=} x_1y_1 + x_2y_2 + \cdots + x_ny_n.$$

Thinking of  $x, y$  as column vectors, this is the same as  $x^T y$ .

### Example

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} = (1 \quad 2 \quad 3) \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} = 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6 = 32.$$

## Properties of the Dot Product

Many usual arithmetic rules hold, as long as you remember you can only dot two vectors together, and that the result is a scalar.

- ▶  $x \cdot y = y \cdot x$
- ▶  $(x + y) \cdot z = x \cdot z + y \cdot z$
- ▶  $(cx) \cdot y = c(x \cdot y)$

Dotting a vector with itself is special:

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1^2 + x_2^2 + \cdots + x_n^2.$$

Hence:

- ▶  $x \cdot x \geq 0$
- ▶  $x \cdot x = 0$  if and only if  $x = 0$ .

**Important:**  $x \cdot y = 0$  does *not* imply  $x = 0$  or  $y = 0$ . For example,  $\begin{pmatrix} 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 0$ .


# The Dot Product and Length

## Definition

The **length** or **norm** of a vector  $x$  in  $\mathbf{R}^n$  is

$$\|x\| = \sqrt{x \cdot x} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}.$$

Why is this a good definition? The Pythagorean theorem!



$$\left\| \begin{pmatrix} 3 \\ 4 \end{pmatrix} \right\| = \sqrt{3^2 + 4^2} = 5$$

## Fact

If  $x$  is a vector and  $c$  is a scalar, then  $\|cx\| = |c| \cdot \|x\|$ .

$$\left\| \begin{pmatrix} 6 \\ 8 \end{pmatrix} \right\| = \left\| 2 \begin{pmatrix} 3 \\ 4 \end{pmatrix} \right\| = 2 \left\| \begin{pmatrix} 3 \\ 4 \end{pmatrix} \right\| = 10$$

## The Dot Product and Distance

### Definition

The **distance** between two points  $x, y$  in  $\mathbf{R}^n$  is


$$\text{dist}(x, y) = \|y - x\|.$$

This is just the length of the vector from  $x$  to  $y$ .

### Example

Let  $x = (1, 2)$  and  $y = (4, 4)$ . Then

$$\text{dist}(x, y) = \|y - x\| = \left\| \begin{pmatrix} 3 \\ 2 \end{pmatrix} \right\| = \sqrt{3^2 + 2^2} = \sqrt{13}.$$



# Unit Vectors

## Definition

A **unit vector** is a vector  $v$  with length  $\|v\| = 1$ .

## Example

The unit coordinate vectors are unit vectors:

$$\|e_1\| = \left\| \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\| = \sqrt{1^2 + 0^2 + 0^2} = 1$$

## Definition

Let  $x$  be a nonzero vector in  $\mathbf{R}^n$ . The **unit vector in the direction of  $x$**  is the vector  $\frac{x}{\|x\|}$ .

This is in fact a unit vector:

$$\text{scalar} \left\| \frac{x}{\|x\|} \right\| = \frac{1}{\|x\|} \|x\| = 1.$$




# Unit Vectors

## Example

### Example

What is the unit vector in the direction of  $x = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ ?

$$u = \frac{x}{\|x\|} = \frac{1}{\sqrt{3^2 + 4^2}} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 3 \\ 4 \end{pmatrix}.$$




# Orthogonality

## Definition

Two vectors  $x, y$  are **orthogonal** or **perpendicular** if  $x \cdot y = 0$ .

*Notation:*  $x \perp y$  means  $x \cdot y = 0$ .

Why is this a good definition? The Pythagorean theorem / law of cosines!



Law of cosines:

$$\|x - y\|^2 = \|x\|^2 + \|y\|^2 - 2\|x\| \|y\| \cos \alpha$$

$$\alpha = 90^\circ \iff \cos \alpha = 0$$

$$x \text{ and } y \text{ are perpendicular} \iff \|x\|^2 + \|y\|^2 = \|x - y\|^2$$

$$\iff x \cdot x + y \cdot y = (x - y) \cdot (x - y)$$

$$\iff x \cdot x + y \cdot y = x \cdot x + y \cdot y - 2x \cdot y$$

$$\iff x \cdot y = 0$$

**Fact:**  $x \perp y \iff \|x - y\|^2 = \|x\|^2 + \|y\|^2$

# Orthogonality

## Example

**Problem:** Find *all* vectors orthogonal to  $v = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$ .

We have to find all vectors  $x$  such that  $x \cdot v = 0$ . This means solving the equation

$$0 = x \cdot v = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = x_1 + x_2 - x_3.$$

The parametric form for the solution is  $x_1 = -x_2 + x_3$ , so the parametric vector form of the general solution is

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$$

For instance,  $\begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \perp \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$  because  $\begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0$ .

# Orthogonality

## Example

**Problem:** Find *all* vectors orthogonal to both  $v = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$  and  $w = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ .

Now we have to solve the system of two homogeneous equations

$$0 = x \cdot v = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = x_1 + x_2 - x_3$$

$$0 = x \cdot w = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = x_1 + x_2 + x_3.$$

In matrix form:

The rows are  $v$  and  $w \rightarrow \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$

The parametric vector form of the solution is

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}.$$

# Orthogonality

## General procedure

**Problem:** Find all vectors orthogonal to some number of vectors  $v_1, v_2, \dots, v_m$  in  $\mathbf{R}^n$ .

This is the same as finding all vectors  $x$  such that

$$0 = v_1^T x = v_2^T x = \dots = v_m^T x.$$

Putting the *row* vectors  $v_1^T, v_2^T, \dots, v_m^T$  into a matrix, this is the same as finding all  $x$  such that

$$\begin{pmatrix} -v_1^T - \\ -v_2^T - \\ \vdots \\ -v_m^T - \end{pmatrix} x = \begin{pmatrix} v_1 \cdot x \\ v_2 \cdot x \\ \vdots \\ v_m \cdot x \end{pmatrix} = 0.$$

### Important

The set of all vectors orthogonal to some vectors  $v_1, v_2, \dots, v_m$  in  $\mathbf{R}^n$  is the *null space* of the  $m \times n$  matrix you get by “turning them sideways and smushing them together:”

$$\begin{pmatrix} -v_1^T - \\ -v_2^T - \\ \vdots \\ -v_m^T - \end{pmatrix}.$$

In particular, this set is a subspace!

## Summary

- ▶ The **dot product** of vectors  $x, y$  in  $\mathbf{R}^n$  is the number  $x^T y$ .
- ▶ The **length** or **norm** of a vector  $x$  in  $\mathbf{R}^n$  is  $\|x\| = \sqrt{x \cdot x}$ .
- ▶ The **distance** between two vectors  $x, y$  in  $\mathbf{R}^n$  is  $\text{dist}(x, y) = \|y - x\|$ .
- ▶ A **unit vector** is a vector  $v$  with length  $\|v\| = 1$ .
- ▶ The **unit vector in the direction of**  $x$  is  $x/\|x\|$ .
- ▶ Two vectors  $x, y$  are **orthogonal** if  $x \cdot y = 0$ .
- ▶ The set of all vectors orthogonal to some vectors  $v_1, v_2, \dots, v_m$  in  $\mathbf{R}^n$  is the null space of the matrix

$$\begin{pmatrix} -v_1^T - \\ -v_2^T - \\ \vdots \\ -v_m^T - \end{pmatrix}.$$