

Math 1553 Worksheet §2.1, §2.2

Solutions

- Write a set of **three** vectors whose span is a **point** in \mathbf{R}^3 .
 - Write a set of **three** different vectors whose span is a **line** in \mathbf{R}^3 .
 - Write a set of **three** different vectors whose span is a **plane** in \mathbf{R}^3 .
 - In each of the above questions, if you use the three vectors form a matrix A , how many pivots does A have?

Solution.

- The span of any three vectors v_1, v_2, v_3 in \mathbf{R}^3 must contain the origin, since

$$0v_1 + 0v_2 + 0v_3 \text{ is automatically the zero vector.}$$

There is only one possibility for this answer: we must choose $v_1 = v_2 = v_3 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$. If our list had contained a nonzero vector, then the span would include that nonzero vector and all scalar multiples of it (including the zero vector).

- Just choose any vector that spans your favorite line, then pick the other vectors to be within that line. For example, choose $v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, $v_2 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$, and $v_3 = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}$, which span the x -axis within \mathbf{R}^3 .

- Similar to above. Just choose any two vectors that span your favorite plane, then pick your third vector to be within that plane. For example, choose $v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, $v_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$, and $v_3 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$. The span of these three vectors is the xy -plane in \mathbf{R}^3 .

- For a) the matrix A has no pivot. For b) the matrix has one pivot. For c) the matrix has two pivots.

- Consider the system of linear equations

$$\begin{aligned}x + 2y &= 7 \\2x + y &= -2 \\-x - y &= 4.\end{aligned}$$

Question: Does this system have a solution? If so, what is the solution set?

- Formulate this question as a question about an augmented matrix.
- Formulate this question as a vector equation.

- c) What does this question mean in terms of spans?
- d) Answer part (c) using the [interactive demo](#).
- e) Answer the question using row reduction.

Solution.

- a) Our question asks whether the augmented matrix below represents a consistent linear system.

$$\left(\begin{array}{cc|c} 1 & 2 & 7 \\ 2 & 1 & -2 \\ -1 & -1 & 4 \end{array} \right)$$

- b) What are the solutions to the following vector equation?

$$x \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ -2 \\ 4 \end{pmatrix}$$

- c) Is $\begin{pmatrix} 7 \\ -2 \\ 4 \end{pmatrix}$ in $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \right\}$?

- d) The picture in the interactive demo shows that b is not in the span of the columns of A , so the system of linear equations is inconsistent.
- e) From part e we already know the system is inconsistent, so here we confirm it using row reduction. Row reducing the matrix in part a yields

$$\left(\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right),$$

so there are no solutions to the system of linear equations.

3. Jameson Locke has challenged you to find a hidden treasure, located at some point (a, b, c) . He has honestly guaranteed you that the treasure can be found by starting at the origin and taking steps in directions given by

$$v_1 = \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} \quad v_2 = \begin{pmatrix} 5 \\ -4 \\ -7 \end{pmatrix} \quad v_3 = \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}.$$

By decoding the message, you have discovered that the first and second coordinates of the treasure's location are (in order) -4 and 3 .

- a) What is the treasure's full location?
- b) Give instructions for how to find the treasure by only moving in the directions given by v_1 , v_2 , and v_3 . Can you do the same to get the treasure by just using v_1 and v_2 ?

Solution.

- a) We translate this problem into linear algebra. Let c be the final entry of the treasure's location. Since Jameson has assured us that we can find the treasure using the three vectors we have been given, our problem is to find c so that

$\begin{pmatrix} -4 \\ 3 \\ c \end{pmatrix}$ is a linear combination of v_1 , v_2 , and v_3 (in other words, find c so that

the treasure's location is in $\text{Span}\{v_1, v_2, v_3\}$). We form an augmented matrix and find when it gives a consistent system.

$$\left(\begin{array}{ccc|c} 1 & 5 & -3 & -4 \\ -1 & -4 & 1 & 3 \\ -2 & -7 & 0 & c \end{array} \right) \xrightarrow[\substack{R_2=R_2+R_1 \\ R_3=R_3+2R_1}]{R_2=R_2+R_1} \left(\begin{array}{ccc|c} 1 & 5 & -3 & -4 \\ 0 & 1 & -2 & -1 \\ 0 & 3 & -6 & c-8 \end{array} \right) \xrightarrow{R_3=R_3-3R_2} \left(\begin{array}{ccc|c} 1 & 5 & -3 & -4 \\ 0 & 1 & -2 & -1 \\ 0 & 0 & 0 & c-5 \end{array} \right).$$

This system will be consistent if and only if the right column is not a pivot column, so we need $c - 5 = 0$, or $c = 5$.

The location of the treasure is $(-4, 3, 5)$.

- b) Getting to the point $(-4, 3, 5)$ using the vectors v_1 , v_2 , and v_3 is equivalent to finding scalars x_1 , x_2 , and x_3 so that

$$\begin{pmatrix} -4 \\ 3 \\ 5 \end{pmatrix} = x_1 \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} + x_2 \begin{pmatrix} 5 \\ -4 \\ -7 \end{pmatrix} + x_3 \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}$$

We can rewrite this as

$$\begin{aligned} x_1 + 5x_2 - 3x_3 &= -4 \\ -x_1 - 4x_2 + x_3 &= 3 \\ -2x_1 - 7x_2 &= 5. \end{aligned}$$

We put the matrix from part (a) into RREF.

$$\left(\begin{array}{ccc|c} 1 & 5 & -3 & -4 \\ 0 & 1 & -2 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{R_1=R_1-5R_2} \left(\begin{array}{ccc|c} 1 & 0 & 7 & 1 \\ 0 & 1 & -2 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Note x_3 is the only free variable, so:

$$x_1 = 1 - 7x_3, \quad x_2 = -1 + 2x_3, \quad x_3 = x_3 \quad (x_3 \text{ real}).$$

Since the system has infinitely many solutions, there are infinitely many ways to get to the treasure. In fact, we can get to the treasure using v_1 and v_2 alone if we wish! If we choose the path corresponding to $x_3 = 0$, then $x_1 = 1$ and $x_2 = -1$, which means that we move 1 unit in the direction of v_1 and -1 unit in the direction of v_2 . In equations:

$$\begin{pmatrix} -4 \\ 3 \\ 5 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} - \begin{pmatrix} 5 \\ -4 \\ -7 \end{pmatrix} + 0 \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}.$$

4. Let $v_1 = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$ $v_2 = \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix}$ $w = \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix}$.

Question: Is w a linear combination of v_1 and v_2 ? In other words, is w in $\text{Span}\{v_1, v_2\}$?

- a) Formulate this question as a vector equation.
- b) Answer the question using the [interactive demo](#).

Solution.

- a) Does the following vector equation have a solution?

$$x \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + y \begin{pmatrix} -2 \\ -3 \\ -1 \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ 8 \end{pmatrix}$$

- b) The demo shows us that w is indeed in the plane spanned by v_1 and v_2 , so we just need to find the coefficients. Using the grid given by the demo, we see $x = 7/2$ and $y = 5/2$.