

### Supplemental problems: §5.2

1. True or false. If the statement is always true, answer true and justify why it is true. Otherwise, answer false and give an example that shows it is false.
  - a) If  $A$  and  $B$  are  $n \times n$  matrices with the same eigenvectors, then  $A$  and  $B$  have the same characteristic polynomial.
  - b) If  $A$  is a  $3 \times 3$  matrix with characteristic polynomial  $-\lambda^3 + \lambda^2 + \lambda$ , then  $A$  is invertible.

#### Solution.

- a) False:  $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  have the same eigenvectors (all nonzero vectors in  $\mathbf{R}^2$ ) but characteristic polynomials  $\lambda^2$  and  $(1 - \lambda)^2$ , respectively.
  - b) False:  $\lambda = 0$  is a root of the characteristic polynomial, so 0 is an eigenvalue, and  $A$  is not invertible.
2. Find all values of  $a$  so that  $\lambda = 1$  an eigenvalue of the matrix  $A$  below.

$$A = \begin{pmatrix} 3 & -1 & 0 & a \\ a & 2 & 0 & 4 \\ 2 & 0 & 1 & -2 \\ 13 & a & -2 & -7 \end{pmatrix}$$

#### Solution.

We need to know which values of  $a$  make the matrix  $A - I_4$  noninvertible. We have

$$A - I_4 = \begin{pmatrix} 2 & -1 & 0 & a \\ a & 1 & 0 & 4 \\ 2 & 0 & 0 & -2 \\ 13 & a & -2 & -8 \end{pmatrix}.$$

We expand cofactors along the third column, then the second column:

$$\begin{aligned} \det(A - I_4) &= 2 \det \begin{pmatrix} 2 & -1 & a \\ a & 1 & 4 \\ 2 & 0 & -2 \end{pmatrix} \\ &= (2)(1) \det \begin{pmatrix} a & 4 \\ 2 & -2 \end{pmatrix} + (2)(1) \det \begin{pmatrix} 2 & a \\ 2 & -2 \end{pmatrix} \\ &= 2(-2a - 8) + 2(-4 - 2a) = -8a - 24. \end{aligned}$$

This is zero if and only if  $a = -3$ .

3. If  $A$  is an  $n \times n$  matrix and  $\det(A) = 2$ , then 2 is an eigenvalue of  $A$ .

#### Solution.

- a) False. For example,  $A = \begin{pmatrix} 4 & 0 \\ 0 & 1/2 \end{pmatrix}$  has  $\det(A) = 2$  but its eigenvalues are 4 and  $\frac{1}{2}$ .

4. Let  $A = \begin{pmatrix} -3 & 0 & -4 \\ 0 & 3 & 0 \\ 6 & 0 & 7 \end{pmatrix}$ .

- a) Find the eigenvalues of  $A$ .  
 b) Find a basis for each eigenspace of  $A$ . Mark your answers clearly.  
 c) Is there a basis of  $\mathbf{R}^3$  that consists of eigenvectors of  $A$ ? Justify your answer.

### Solution.

- a) We solve  $0 = \det(A - \lambda I)$ .

$$\begin{aligned} 0 &= \det \begin{pmatrix} -3-\lambda & 0 & -4 \\ 0 & 3-\lambda & 0 \\ 6 & 0 & 7-\lambda \end{pmatrix} = (3-\lambda)(-1)^4 \det \begin{pmatrix} -3-\lambda & -4 \\ 6 & 7-\lambda \end{pmatrix} \\ &= (3-\lambda)((-3-\lambda)(7-\lambda) + 24) = (3-\lambda)(\lambda^2 - 4\lambda - 21 + 24) \\ &= (3-\lambda)(\lambda^2 - 4\lambda + 3) = (3-\lambda)(\lambda-3)(\lambda-1) \end{aligned}$$

So  $\lambda = 1$  and  $\lambda = 3$  are the eigenvalues.

$$\underline{\lambda = 1}: (A - I \mid 0) = \left( \begin{array}{ccc|c} -4 & 0 & -4 & 0 \\ 0 & 2 & 0 & 0 \\ 6 & 0 & 6 & 0 \end{array} \right) \xrightarrow[\text{then } R_1 = -R_1/4]{R_3 = R_3 + \frac{3}{2}R_1} \left( \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \text{ with so-}$$

lution  $x_1 = -x_3$ ,  $x_2 = 0$ ,  $x_3 = x_3$ . The 1-eigenspace has basis  $\left\{ \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\}$ .

$\lambda = 3$ :

$$(A - 3I \mid 0) = \left( \begin{array}{ccc|c} -6 & 0 & -4 & 0 \\ 0 & 0 & 0 & 0 \\ 6 & 0 & 4 & 0 \end{array} \right) \xrightarrow[\text{then } R_1 = -R_1/6]{R_3 = R_3 + R_1} \left( \begin{array}{ccc|c} 1 & 0 & \frac{2}{3} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

with solution  $x_1 = -\frac{2}{3}x_3$ ,  $x_2 = x_2$ ,  $x_3 = x_3$ . The 3-eigenspace has basis  $\left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2/3 \\ 0 \\ 1 \end{pmatrix} \right\}$ .

- b) Yes. The eigenvectors that we have found form a basis of  $\mathbf{R}^3$ . One step of row-reduction shows that the three eigenvectors in  $\mathbf{R}^3$  below are linearly independent, and are therefore a basis of  $\mathbf{R}^3$  by the Basis Theorem.

$$\left\{ \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2/3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

**Supplemental problems: §5.4**

1. True or false. Answer true if the statement is always true. Otherwise, answer false.
- If  $A$  is an invertible matrix and  $A$  is diagonalizable, then  $A^{-1}$  is diagonalizable.
  - A diagonalizable  $n \times n$  matrix admits  $n$  linearly independent eigenvectors.
  - If  $A$  is diagonalizable, then  $A$  has  $n$  distinct eigenvalues.

**Solution.**

- True. If  $A = PDP^{-1}$  and  $A$  is invertible then its eigenvalues are all nonzero, so the diagonal entries of  $D$  are nonzero and thus  $D$  is invertible (pivot in every diagonal position). Thus,  $A^{-1} = (PDP^{-1})^{-1} = (P^{-1})^{-1}D^{-1}P^{-1} = PD^{-1}P^{-1}$ .
  - True. By the Diagonalization Theorem, an  $n \times n$  matrix is diagonalizable *if and only if* it admits  $n$  linearly independent eigenvectors.
  - False. For instance,  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is diagonal but has only one eigenvalue.
2. Give examples of  $2 \times 2$  matrices with the following properties. Justify your answers.
- A matrix  $A$  which is invertible and diagonalizable.
  - A matrix  $B$  which is invertible but not diagonalizable.
  - A matrix  $C$  which is not invertible but is diagonalizable.
  - A matrix  $D$  which is neither invertible nor diagonalizable.

**Solution.**

- a) We can take any diagonal matrix with nonzero diagonal entries:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

- b) A shear has only one eigenvalue  $\lambda = 1$ . The associated eigenspace is the  $x$ -axis, so there do not exist two linearly independent eigenvectors. Hence it is not diagonalizable.

$$B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

- c) We can take any diagonal matrix with some zero diagonal entries:

$$C = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

- d) Such a matrix can only have the eigenvalue zero — otherwise it would have two eigenvalues, hence be diagonalizable. Thus the characteristic polynomial

is  $f(\lambda) = \lambda^2$ . Here is a matrix with trace and determinant zero, whose zero-eigenspace (i.e., null space) is not all of  $\mathbf{R}^2$ :

$$D = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

3.  $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}.$

- a) Find the eigenvalues of  $A$ , and find a basis for each eigenspace.
- b) Is  $A$  diagonalizable? If your answer is yes, find a diagonal matrix  $D$  and an invertible matrix  $C$  so that  $A = CDC^{-1}$ . If your answer is no, justify why  $A$  is not diagonalizable.

### Solution.

a) We solve  $0 = \det(A - \lambda I)$ .

$$\begin{aligned} 0 &= \det \begin{pmatrix} 2-\lambda & 3 & 1 \\ 3 & 2-\lambda & 4 \\ 0 & 0 & -1-\lambda \end{pmatrix} = (-1-\lambda)(-1)^6 \det \begin{pmatrix} 2-\lambda & 3 \\ 3 & 2-\lambda \end{pmatrix} = (-1-\lambda)((2-\lambda)^2 - 9) \\ &= (-1-\lambda)(\lambda^2 - 4\lambda - 5) = -(\lambda+1)^2(\lambda-5). \end{aligned}$$

So  $\lambda = -1$  and  $\lambda = 5$  are the eigenvalues.

$$\underline{\lambda = -1}: (A + I | 0) = \left( \begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 3 & 3 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{R_2=R_2-R_1} \left( \begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow[\text{then } R_1=R_1/3]{R_1=R_1-R_2}$$

$$\left( \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right), \text{ with solution } x_1 = -x_2, x_2 = x_2, x_3 = 0. \text{ The } (-1)\text{-eigenspace}$$

$$\text{has basis } \left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

$\lambda = 5$ :

$$(A - 5I | 0) = \left( \begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 3 & -3 & 4 & 0 \\ 0 & 0 & -6 & 0 \end{array} \right) \xrightarrow[\begin{smallmatrix} R_2=R_2+R_1 \\ R_3=R_3/(-6) \end{smallmatrix}]{R_2=R_2+R_1} \left( \begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right) \xrightarrow[\text{then } R_2 \leftrightarrow R_3, R_1/(-3)]{R_1=R_1-R_3, R_2=R_2-5R_3} \left( \begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$$

$$\text{with solution } x_1 = x_2, x_2 = x_2, x_3 = 0. \text{ The } 5\text{-eigenspace has basis } \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

- b)  $A$  is a  $3 \times 3$  matrix that only admits 2 linearly independent eigenvectors, so  $A$  is not diagonalizable.

4. Let  $A = \begin{pmatrix} 8 & 36 & 62 \\ -6 & -34 & -62 \\ 3 & 18 & 33 \end{pmatrix}$ .

The characteristic polynomial for  $A$  is  $-\lambda^3 + 7\lambda^2 - 16\lambda + 12$ , and  $\lambda - 3$  is a factor. Decide if  $A$  is diagonalizable. If it is, find an invertible matrix  $C$  and a diagonal matrix  $D$  such that  $A = CDC^{-1}$ .

**Solution.**

By polynomial division,

$$\frac{-\lambda^3 + 7\lambda^2 - 16\lambda + 12}{\lambda - 3} = -\lambda^2 + 4\lambda - 4 = -(\lambda - 2)^2.$$

Thus, the characteristic poly factors as  $-(\lambda - 3)(\lambda - 2)^2$ , so the eigenvalues are  $\lambda_1 = 3$  and  $\lambda_2 = 2$ .

For  $\lambda_1 = 3$ , we row-reduce  $A - 3I$ :

$$\begin{pmatrix} 5 & 36 & 62 \\ -6 & -37 & -62 \\ 3 & 18 & 30 \end{pmatrix} \xrightarrow[\text{(New } R_1)/3]{R_1 \leftrightarrow R_3} \begin{pmatrix} 1 & 6 & 10 \\ -6 & -37 & -62 \\ 5 & 36 & 62 \end{pmatrix} \xrightarrow[R_3 = R_3 - 5R_1]{R_2 = R_2 + 6R_1} \begin{pmatrix} 1 & 6 & 10 \\ 0 & -1 & -2 \\ 0 & 6 & 12 \end{pmatrix}$$

$$\xrightarrow[\text{then } R_2 = -R_2]{R_3 = R_3 + 6R_2} \begin{pmatrix} 1 & 6 & 10 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \xrightarrow{R_1 = R_1 - 6R_2} \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore, the solutions to  $(A - 3I \mid 0)$  are  $x_1 = 2x_3$ ,  $x_2 = -2x_3$ ,  $x_3 = x_3$ .

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2x_3 \\ -2x_3 \\ x_3 \end{pmatrix} = x_3 \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}. \quad \text{The 3-eigenspace has basis } \left\{ \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} \right\}.$$

For  $\lambda_2 = 2$ , we row-reduce  $A - 2I$ :

$$\begin{pmatrix} 6 & 36 & 62 \\ -6 & -36 & -62 \\ 3 & 18 & 31 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 6 & \frac{31}{3} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The solutions to  $(A - 2I \mid 0)$  are  $x_1 = -6x_2 - \frac{31}{3}x_3$ ,  $x_2 = x_2$ ,  $x_3 = x_3$ .

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -6x_2 - \frac{31}{3}x_3 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} -6 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -\frac{31}{3} \\ 0 \\ 1 \end{pmatrix}.$$

The 2-eigenspace has basis  $\left\{ \begin{pmatrix} -6 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -\frac{31}{3} \\ 0 \\ 1 \end{pmatrix} \right\}$ .

Therefore,  $A = CDC^{-1}$  where

$$C = \begin{pmatrix} 2 & -6 & -\frac{31}{3} \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Note that we arranged the eigenvectors in  $C$  in order of the eigenvalues 3, 2, 2, so we had to put the diagonals of  $D$  in the same order.

5. Which of the following  $3 \times 3$  matrices are necessarily diagonalizable over the real numbers? (Circle all that apply.)
1. A matrix with three distinct real eigenvalues.
  2. A matrix with one real eigenvalue.
  3. A matrix with a real eigenvalue  $\lambda$  of algebraic multiplicity 2, such that the  $\lambda$ -eigenspace has dimension 2.
  4. A matrix with a real eigenvalue  $\lambda$  such that the  $\lambda$ -eigenspace has dimension 2.

**Solution.**

The matrices in 1 and 3 are diagonalizable. A matrix with three distinct real eigenvalues automatically admits three linearly independent eigenvectors. If a matrix  $A$  has a real eigenvalue  $\lambda_1$  of algebraic multiplicity 2, then it has another real eigenvalue  $\lambda_2$  of algebraic multiplicity 1. The two eigenspaces provide three linearly independent eigenvectors.

The matrices in 2 and 4 need not be diagonalizable.

6. Suppose a  $2 \times 2$  matrix  $A$  has eigenvalue  $\lambda_1 = -2$  with eigenvector  $v_1 = \begin{pmatrix} 3/2 \\ 1 \end{pmatrix}$ , and eigenvalue  $\lambda_2 = -1$  with eigenvector  $v_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ .
- a) Find  $A$ .
  - b) Find  $A^{100}$ .

**Solution.**

a) We have  $A = CDC^{-1}$  where

$$C = \begin{pmatrix} 3/2 & 1 \\ 1 & -1 \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} -2 & 0 \\ 0 & -1 \end{pmatrix}.$$


$$\text{We compute } C^{-1} = \frac{1}{-5/2} \begin{pmatrix} -1 & -1 \\ -1 & 3/2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 2 & 2 \\ 2 & -3 \end{pmatrix}.$$

$$A = CDC^{-1} = \frac{1}{5} \begin{pmatrix} 3/2 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -2 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 2 & 2 \\ 2 & -3 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} -8 & -3 \\ -2 & -7 \end{pmatrix}.$$

b)

$$\begin{aligned}
 A^{100} &= CD^{100}C^{-1} = \frac{1}{5} \begin{pmatrix} 3/2 & 1 \\ 1 & -1 \end{pmatrix} \cdot D^{100} \begin{pmatrix} 2 & 2 \\ 2 & -3 \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 3/2 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 2^{100} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 2 \\ 2 & -3 \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 3/2 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 2 \cdot 2^{100} & 2 \cdot 2^{100} \\ 2 & -3 \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 3 \cdot 2^{100} + 2 & 3 \cdot 2^{100} - 3 \\ 2^{101} - 2 & 2^{101} + 3 \end{pmatrix}.
 \end{aligned}$$

7. Suppose that  $A = C \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix} C^{-1}$ , where  $C$  has columns  $v_1$  and  $v_2$ . Given  $x$  and  $y$  in the picture below, draw the vectors  $Ax$  and  $Ay$ .

**Solution.**

$A$  does the same thing as  $D = \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix}$ , but in the  $v_1, v_2$ -coordinate system. Since  $D$  scales the first coordinate by  $1/2$  and the second coordinate by  $-1$ , hence  $A$  scales the  $v_1$ -coordinate by  $1/2$  and the  $v_2$ -coordinate by  $-1$ .