

Math 1553 Worksheet §6.1 - §6.5

Solutions

1. True/False. Justify your answer.

- (1) If u is in subspace W , and u is also in W^\perp , then $u = 0$.
- (2) If y is in a subspace W , the orthogonal projection of y onto W^\perp is 0 .
- (3) If x is orthogonal to v and w , then x is also orthogonal to $v - w$.

Solution.

- (1) TRUE: Such a vector u would be orthogonal to itself, so $u \cdot u = \|u\|^2 = 0$. Therefore, u has length 0, so $u = 0$.
- (2) TRUE: y is in W , so $y \perp W^\perp$. Its orthogonal projection onto W is y and orthogonal projection onto W^\perp is 0 . In fact y has orthogonal decomposition $y = y + 0$, where y is in W and 0 is in W^\perp .
- (3) TRUE: By properties of the dot product, if x is orthogonal to v and w then x is orthogonal to everything in $\text{Span}\{v, w\}$ (which includes $v - w$).

2. a) Find the standard matrix B for proj_W , where $W = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} \right\}$.

b) What are the eigenvalues of B ? Is B diagonalizable?

c) Let $x = \begin{pmatrix} 3 \\ 0 \\ 9 \end{pmatrix}$. Find the orthogonal decomposition of x with respect to W .

In other words, find x_W in W and x_{W^\perp} in W^\perp so that $x = x_W + x_{W^\perp}$.

Solution.

a) We use the formula $B = \frac{1}{u \cdot u} uu^T$ where $u = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$ (this is the formula

$B = A(A^T A)^{-1} A^T$ when “ A ” is just the single vector u).

$$B = \frac{1}{1(1) + 1(1) + (-1)(-1)} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} (1 \ 1 \ -1) = \frac{1}{3} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix}$$

$$\implies B = \frac{1}{3} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix}.$$

b) $Bx = x$ for every x in W , and $Bx = 0$ for every x in W^\perp , so B has two eigenvalues: $\lambda_1 = 1$ with algebraic and geometric multiplicity 1, $\lambda_2 = 0$ with algebraic and geometric multiplicity 2. Therefore, B is diagonalizable. We can actually compute the diagonalization of B (we’re not asked in the question). Here

$v_1 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$ is an eigenvector for $\lambda_1 = 1$, whereas $v_2 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$ and $v_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ are linearly independent vectors that are orthogonal to v_1 , so they span the eigenspace for $\lambda_2 = 0$. Therefore

$$B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ -1 & 0 & 1 \end{pmatrix}^{-1}$$

c) It follows that

$$x_W = \frac{1}{3} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 0 \\ 9 \end{pmatrix} = \begin{pmatrix} -2 \\ -2 \\ 2 \end{pmatrix}.$$

Hence, as

$$x_{W^\perp} = x - x_W,$$

we have that

$$x_{W^\perp} = \begin{pmatrix} 3 \\ 0 \\ 9 \end{pmatrix} - \begin{pmatrix} -2 \\ -2 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 2 \\ 7 \end{pmatrix}.$$

$$\text{Thus, } x = \begin{pmatrix} -2 \\ -2 \\ 2 \end{pmatrix} + \begin{pmatrix} 5 \\ 2 \\ 7 \end{pmatrix}.$$

3. Use least-squares to find the best fit line $y = Ax + B$ through the points $(0, 0)$, $(1, 8)$, $(3, 8)$, and $(4, 20)$.

Solution.

We want to find a least squares solution to the system of linear equations

$$\begin{aligned} 0 &= A(0) + B \\ 8 &= A(1) + B \\ 8 &= A(3) + B \\ 20 &= A(4) + B \end{aligned} \iff \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 3 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ 8 \\ 20 \end{pmatrix}.$$

We compute

$$\begin{pmatrix} 0 & 1 & 3 & 4 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 26 & 8 \\ 8 & 4 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 & 3 & 4 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 8 \\ 8 \\ 20 \end{pmatrix} = \begin{pmatrix} 112 \\ 36 \end{pmatrix}$$

$$\left(\begin{array}{cc|cc} 26 & 8 & 112 & \\ 8 & 4 & 36 & \end{array} \right) \xrightarrow{\text{rref}} \left(\begin{array}{cc|cc} 1 & 0 & 4 & \\ 0 & 1 & 1 & \end{array} \right).$$

Hence the least squares solution is $A = 4$ and $B = 1$, so the best fit line is $y = 4x + 1$.