

## Math 1553 Worksheet: Sections 5.1-5.2

1. True or false: If  $v_1$  and  $v_2$  are linearly independent eigenvectors of an  $n \times n$  matrix  $A$ , then they must correspond to different eigenvalues.


### Solution.

False. For example, if  $A = I_2$  then  $e_1$  and  $e_2$  are linearly independent eigenvectors both corresponding to the eigenvalue  $\lambda = 1$ .

2. In what follows,  $T$  is a linear transformation with matrix  $A$ . Find the eigenvectors and eigenvalues of  $A$  without doing any matrix calculations. (Draw a picture!)
- $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  that projects vectors onto the  $xz$ -plane in  $\mathbb{R}^3$ .
  - $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  that reflects vectors over the line  $y = 2x$  in  $\mathbb{R}^2$ .

### Solution.

- a) We draw the  $xz$ -plane below.




$T(x, y, z) = (x, 0, z)$ , so  $T$  fixes every vector in the  $xz$ -plane and destroys every vector of the form  $(0, a, 0)$  with  $a$  real. Therefore,  $\lambda = 1$  and  $\lambda = 0$  are eigenvalues and in fact they are the only eigenvalues since their combined eigenvectors span all of  $\mathbb{R}^3$ .

The eigenvectors for  $\lambda = 1$  are all vectors of the form  $\begin{pmatrix} x \\ 0 \\ z \end{pmatrix}$  where at least one of  $x$  and  $z$  is nonzero, and the eigenvectors for  $\lambda = 0$  are all vectors of the form  $\begin{pmatrix} 0 \\ y \\ 0 \end{pmatrix}$  where  $y \neq 0$ . In other words:

The 1-eigenspace consists of all vectors in  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$ , while the 0-eigenspace consists of all vectors in  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}$

b) Here is the picture:



$T$  fixes every vector along the line  $y = 2x$ , so  $\lambda = 1$  is an eigenvalue and its eigenvectors are all vectors  $\begin{pmatrix} t \\ 2t \end{pmatrix}$  where  $t \neq 0$ .

$T$  flips every vector along the line perpendicular to  $y = 2x$ , which is  $y = -\frac{1}{2}x$  (for example,  $T(-2, 1) = (2, -1)$ ). Therefore,  $\lambda = -1$  is an eigenvalue and its eigenvectors are all vectors of the form  $\begin{pmatrix} s \\ -\frac{1}{2}s \end{pmatrix}$  where  $s \neq 0$ .

3. True or False: Suppose  $A = \begin{pmatrix} 3 & 0 & 0 \\ 5 & 1 & 0 \\ -10 & 4 & 7 \end{pmatrix}$ . Then the characteristic polynomial of  $A$  is

$$\det(A - \lambda I) = (3 - \lambda)(1 - \lambda)(7 - \lambda).$$

**Solution.**

Yes. Since  $A - \lambda I$  is triangular, its determinant is the product of its diagonal entries.

4. Find the eigenvalues and a basis for each eigenspace of  $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}$ .

**Solution.**

We solve  $0 = \det(A - \lambda I)$ .

$$\begin{aligned} 0 &= \det \begin{pmatrix} 2-\lambda & 3 & 1 \\ 3 & 2-\lambda & 4 \\ 0 & 0 & -1-\lambda \end{pmatrix} = (-1-\lambda)(-1)^6 \det \begin{pmatrix} 2-\lambda & 3 \\ 3 & 2-\lambda \end{pmatrix} = (-1-\lambda)((2-\lambda)^2 - 9) \\ &= (-1-\lambda)(\lambda^2 - 4\lambda - 5) = -(\lambda+1)^2(\lambda-5). \end{aligned}$$

So  $\lambda = -1$  and  $\lambda = 5$  are the eigenvalues.

$$\underline{\lambda = -1}: (A + I | 0) = \left( \begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 3 & 3 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{R_2=R_2-R_1} \left( \begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{\substack{R_1=R_1-R_2 \\ \text{then } R_1=R_1/3}} \left( \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$$

with solution  $x_1 = -x_2$ ,  $x_2 = x_2$ ,  $x_3 = 0$ . The  $(-1)$ -eigenspace has basis  $\left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \right\}$ .

$\lambda = 5$ :

$$(A - 5I | 0) = \left( \begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 3 & -3 & 4 & 0 \\ 0 & 0 & -6 & 0 \end{array} \right) \xrightarrow{\substack{R_2=R_2+R_1 \\ R_3=R_3/(-6)}} \left( \begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right) \xrightarrow{\substack{R_1=R_1-R_3, R_2=R_2-5R_3 \\ \text{then } R_2 \leftrightarrow R_3, R_1/(-3)}} \left( \begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$$

with solution  $x_1 = x_2$ ,  $x_2 = x_2$ ,  $x_3 = 0$ . The 5-eigenspace has basis  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}$ .