

(Reduced) Row Echelon Form

1.2 Review

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.
2. Each leading nonzero entry of a row is to the *right* of the leading entry of the row above.
3. Below a leading entry of a row, all entries are *zero*.

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Row echelon form:

$$\begin{pmatrix} \boxed{\star} & \star & \star & \star & \star \\ 0 & \boxed{\star} & \star & \star & \star \\ 0 & 0 & 0 & \boxed{\star} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Reduced row echelon form:

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$\boxed{\star} = \text{pivots}$

1.2: Row Reduction

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction** or **Gaussian elimination**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form. (Assuming you only do the three legal row operations.) (And you don't make any arithmetic errors.)

Maybe you can figure out why it's true!

Reinforcing 1.2: Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries below this 1 are 0.

Step 2a Swap the 2nd row with a lower one so that the leftmost nonzero entry is in 2nd row.

Step 2b Scale 2nd row so that its leading entry is equal to 1.

Step 2c Use row replacement so all entries below this 1 are 0.

Step 3a Swap the 3rd row with a lower one so that the leftmost nonzero entry is in 3rd row.

etc.

Last Step Use row replacement to clear all entries above the pivots, starting with the last pivot (to make life easier).

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

[animated]

Row Reduction

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -5 & -10 & -20 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

Note: Step 2 never messes up the first (nonzero) column of the matrix, because it looks like this:

"Active" row $\rightarrow$ $\left(\begin{array}{ccc|c} 1 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{array} \right)$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Row Reduction

Example, continued

Success! The reduced row echelon form is

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right) \implies \begin{cases} x & = & 1 \\ y & = & -2 \\ z & = & 3 \end{cases}$$

Recap

Get a 1 here

$$\begin{pmatrix} \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \end{pmatrix}$$

Get a 1 here

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \end{pmatrix}$$

(maybe these are already zero)

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & \star \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Get a 1 here

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & \star \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Matrix is in REF

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Clear up

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Clear up

$$\begin{pmatrix} 1 & \star & \star & 0 \\ 0 & 1 & \star & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Matrix is in RREF

$$\begin{pmatrix} 1 & 0 & \star & 0 \\ 0 & 1 & \star & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Profit?

Row Reduction

Another example

The linear system

$$2x + 10y = -1$$

$$3x + 15y = 2$$

gives rise to the matrix

Let's row reduce it: [\[interactive row reducer\]](#)

$$\left(\begin{array}{cc|c} 2 & 10 & -1 \\ 3 & 15 & 2 \end{array} \right)$$

The row reduced matrix

$$\left(\begin{array}{cc|c} 1 & 5 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

corresponds to the
inconsistent system

$$\begin{aligned} x + 5y &= 0 \\ 0 &= 1. \end{aligned}$$

Question

What does an augmented matrix in reduced row echelon form look like, if its system of linear equations is inconsistent?

An augmented matrix corresponds to an inconsistent system of equations if and only if *the last* (i.e., the augmented) *column is a pivot column*.

Section 1.3

Parametric Form

Example 1

How do we solve a system of linear equations if the row reduced matrix has a column without a pivot? Let's do an example.

$$\begin{array}{rcl} 2x + y + 12z & = & 1 \\ x + 2y + 9z & = & -1 \end{array} \quad \text{gives rise to the matrix} \quad \left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right).$$

Let's row reduce it: [\[interactive row reducer\]](#)

$$\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$$

The row reduced matrix

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 1 \\ 0 & 1 & 2 & -1 \end{array} \right)$$

corresponds to the
linear system

$$\begin{cases} x + 5z = 1 \\ y + 2z = -1 \end{cases}$$

Example 1

Continued

Definition

Consider a *consistent* linear system of equations in the variables $x_1, \dots, x_n$. Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example, z was free because the reduced row echelon form matrix was

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

Example 2

Suppose the reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

Example 3

Solve the system of linear equations in x_1, x_2, x_3, x_4 :

$$x_1 + 5x_3 = 0$$

$$x_4 = 0$$

So the associated matrix is:



$$\left(\begin{array}{cccc|c} 1 & 0 & 5 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right)$$

Free variables

Geometry, looking ahead!

If we have a consistent system of linear equations, with n variables and k free variables, then the set of solutions is a k -dimensional object in $\mathbf{R}^n$. We will make this precise later, but it is worth thinking about now.

Why does this make sense?



A linear system has 4 variables and 3 equations.

What are the possible solution sets?

- (a) point
- (b) two points
- (c) line
- (d) plane
- (e) 3-dimensional object
- (f) 4-dimensional object

Yet Another Example

The linear system

$$x + y + z = 1 \quad \text{has matrix form}$$

This is in reduced row echelon form. The free variables are y and z . The parametric form of the general solution is

Rearranging:

$$(x, y, z) = (1 - y - z, y, z),$$

where y and z are arbitrary real numbers.

[interactive]

Trichotomy

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are zero solutions, i.e. the solution set is *empty*. Picture:

$$\left(\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

2. Every column except the last column is a pivot column.

In this case, the system has a *unique solution*. Picture:

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & \star \\ 0 & 1 & 0 & \star \\ 0 & 0 & 1 & \star \end{array} \right)$$

3. The last column is not a pivot column, and some other column isn't either.

In this case, the system has *infinitely many* solutions, corresponding to the infinitely many possible values of the free variable(s). Picture:

$$\left(\begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$

Summary

- ▶ **Row reduction** is an algorithm for solving a system of linear equations represented by an augmented matrix.
- ▶ The goal of row reduction is to put a matrix into **(reduced) row echelon form**, which is the “solved” version of the matrix.
- ▶ An augmented matrix corresponds to an inconsistent system if and only if there is a pivot in the augmented column.
- ▶ Columns without pivots in the RREF of a matrix correspond to **free variables**. You can assign any value you want to the free variables.
- ▶ We can tell whether a linear system has zero, one, or infinitely many solutions using the RREF of the augmented matrix.