

Math 1553 Worksheet §3.2, 3.3

Solutions

1. Which of the following statements are true? Justify your answer.

a) Let A be a 3×3 matrix, such that $Ax = \begin{pmatrix} 1 \\ 5 \\ 2 \end{pmatrix}$ has a unique solution. Then,

$Ax = \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix}$ also has a unique solution.

b) Let A be a 3×4 matrix. Then, the transformation whose standard matrix is A cannot be onto.

Solution.

a) True. If $Ax = b$ has a unique solution (regardless of what b is), then so does $Ax = 0$, so the transformation whose standard matrix is A is one-to-one. Therefore, A has a pivot in each column (3 pivots). Since A is 3×3 , this means A also has a pivot each row, so for every vector b' in \mathbf{R}^3 , the system $Ax = b'$ is consistent and has unique solution.

b) False. The matrix $A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$ is such matrix, since it has pivots in every row.

2. Which of the following transformations T are onto? Which are one-to-one? If the transformation is not onto, find a vector not in the range. If the transformation is not one-to-one, find two vectors with the same image.
- Counterclockwise rotation by 32° in \mathbf{R}^2 .
 - The transformation $T : \mathbf{R}^3 \rightarrow \mathbf{R}^2$ defined by $T(x, y, z) = (z, x)$.
 - The matrix transformation with standard matrix $A = \begin{pmatrix} 1 & 6 \\ -1 & 2 \\ 2 & -1 \end{pmatrix}$.

Solution.

- This is both one-to-one and onto. If v is any vector in \mathbf{R}^2 , then there is one and only one vector w such that $T(w) = v$, namely, the vector w that is rotated 32° clockwise from v .
- This is onto. If (a, b) is any vector in the codomain \mathbf{R}^2 , then $(a, b) = T(b, 0, a)$, so (a, b) is in the range. It is not one-to-one though: indeed, $T(0, 0, 0) = (0, 0) = T(0, 1, 0)$. Alternatively, we could have observed that T is a matrix transformation and examined its matrix A : $T(x) = Ax$ for

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}.$$

Since A has a pivot in every row but not every column, T is onto but not one-to-one.

- The transformation T with matrix A is onto if and only if A has a pivot in every row, and it is one-to-one if and only if A has a pivot in every column. So we row reduce:

$$A = \begin{pmatrix} 1 & 6 \\ -1 & 2 \\ 2 & -1 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

This has a pivot in every column, so T is one-to-one. It does not have a pivot in every row, so it is not onto. To find a specific vector b in \mathbf{R}^3 which is not in the image of T , we have to find a $b = (b_1, b_2, b_3)$ such that the matrix equation $Ax = b$ is inconsistent. We row reduce again:

$$(A | b) = \left(\begin{array}{cc|c} 1 & 6 & b_1 \\ -1 & 2 & b_2 \\ 2 & -1 & b_3 \end{array} \right) \xrightarrow{\text{rref}} \left(\begin{array}{cc|c} 1 & 0 & \text{don't care} \\ 0 & 1 & \text{don't care} \\ 0 & 0 & -3b_1 + 13b_2 + 8b_3 \end{array} \right).$$

Hence any b_1, b_2, b_3 for which $-3b_1 + 13b_2 + 8b_3 \neq 0$ will make the equation $Ax = b$ inconsistent. For instance, $b = (1, 0, 0)$ is not in the range of T .

3. The second little pig has decided to build his house out of sticks. The big bad wolf finds the pig's house and blows it down so that the house is rotated by an angle of 45° in a counterclockwise direction about the z -axis (look downward onto the xy -plane the way we usually picture the plane as \mathbf{R}^2), and then projected onto the xy -plane. Find the standard matrix A for the transformation T caused by the wolf.

Solution.

To compute the matrix for T , we have to compute $T(e_1)$, $T(e_2)$, and $T(e_3)$. To see the picture, let's put ourselves above the xy -plane (with the usual orientation of the x and y axes in the xy -plane), looking downward. For e_1 and e_2 , it is as if we are applying $\begin{pmatrix} \cos(45^\circ) & -\sin(45^\circ) \\ \sin(45^\circ) & \cos(45^\circ) \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$ to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, then putting a zero in the z -coordinate each time. We find

$$T(e_1) = T \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad T(e_2) = T \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}.$$

Rotating e_3 around the z -axis does nothing, and projecting onto the xy -plane sends it to zero, so $T(e_3) = 0$. Therefore, the matrix for T is

$$A = \begin{pmatrix} | & | & | \\ T(e_1) & T(e_2) & T(e_3) \\ | & | & | \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$