

## Math 1553 Worksheet §5.2 - §5.4

### Solutions

1. Answer yes, no, or maybe. Justify your answers. In each case,  $A$  is a matrix whose entries are real numbers.

a) Suppose  $A = \begin{pmatrix} 3 & 0 & 0 \\ 5 & 1 & 0 \\ -10 & 4 & 7 \end{pmatrix}$ . Then the characteristic polynomial of  $A$  is

$$\det(A - \lambda I) = (3 - \lambda)(1 - \lambda)(7 - \lambda).$$

- b) If  $A$  is a  $3 \times 3$  matrix with characteristic polynomial  $-\lambda(\lambda - 5)^2$ , then the 5-eigenspace is 2-dimensional.

- c) If  $A$  is an invertible  $2 \times 2$  matrix, then  $A$  is diagonalizable.

### Solution.

- a) Yes. Since  $A - \lambda I$  is triangular, its determinant is the product of its diagonal entries.

- b) Maybe. The geometric multiplicity of  $\lambda = 5$  can be 1 or 2. For example, the


matrix  $\begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 0 \end{pmatrix}$  has a 5-eigenspace which is 2-dimensional, whereas the

matrix  $\begin{pmatrix} 5 & 1 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 0 \end{pmatrix}$  has a 5-eigenspace which is 1-dimensional. Both matrices

have characteristic polynomial  $-\lambda(5 - \lambda)^2$ .

- c) Maybe. The identity matrix is invertible and diagonalizable, but the matrix  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  is invertible but not diagonalizable.

2. The eigenspaces of some  $2 \times 2$  matrix  $A$  are drawn below. Write an invertible matrix  $C$  and a diagonal matrix  $D$  so that  $A = CDC^{-1}$ . Can you find another pair of  $C$  and  $D$  so that  $A = CDC^{-1}$ ?



### Solution.

We choose  $D$  to be a diagonal matrix whose entries are the eigenvalues of  $A$ , and  $C$  a matrix whose columns are corresponding eigenvectors (written in the same order).

The eigenvalues of  $A$  are  $\lambda_1 = -1$  and  $\lambda_2 = -2$ .

The  $(-1)$ -eigenspace is spanned by  $v_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ .

The  $(-2)$ -eigenspace is spanned by  $v_2 = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$ .

Therefore, we can choose  $C = (v_1 \ v_2) = \begin{pmatrix} 1 & 3 \\ -1 & 2 \end{pmatrix}$  and  $D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -2 \end{pmatrix}$ .

There are many possibilities for  $C$  and  $D$ .

For example, since  $\text{Span} \left\{ \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}$ , we could have chosen

$v_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$  instead, so that

$$C = \begin{pmatrix} -1 & 3 \\ 1 & 2 \end{pmatrix}, \quad D = \begin{pmatrix} -1 & 0 \\ 0 & -2 \end{pmatrix}.$$

Alternatively, we could have rearranged the order of the diagonal entries of  $D$  and taken care to use the corresponding order in the columns of  $C$ :

$$C = \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix}, \quad D = \begin{pmatrix} -2 & 0 \\ 0 & -1 \end{pmatrix}.$$

Regardless, if you write any correct answers for  $C$  and  $D$  and go the extra step of carrying out the computation, you will obtain

$$A = CDC^{-1} = -\frac{1}{5} \begin{pmatrix} 8 & 3 \\ 2 & 7 \end{pmatrix}.$$

3. Suppose  $A$  is a  $2 \times 2$  matrix satisfying

$$A \begin{pmatrix} -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \end{pmatrix}, \quad A \begin{pmatrix} -2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

a) Diagonalize  $A$  by finding  $2 \times 2$  matrices  $C$  and  $D$  (with  $D$  diagonal) so that  $A = CDC^{-1}$ .

b) Find  $A^{17}$ .

### Solution.

a) From the information given,  $\lambda_1 = -2$  is an eigenvalue for  $A$  with corresponding eigenvector  $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ , and  $\lambda_2 = 0$  is an eigenvalue with eigenvector  $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$ .

By the Diagonalization Theorem,  $A = CDC^{-1}$  where

$$C = \begin{pmatrix} -1 & -2 \\ 1 & 3 \end{pmatrix}, \quad D = \begin{pmatrix} -2 & 0 \\ 0 & 0 \end{pmatrix}.$$

b) We find  $C^{-1} = \frac{1}{-3+2} \begin{pmatrix} 3 & 2 \\ -1 & -1 \end{pmatrix} = \begin{pmatrix} -3 & -2 \\ 1 & 1 \end{pmatrix}$ .

$$\begin{aligned} A^{17} &= CD^{17}C^{-1} = \begin{pmatrix} -1 & -2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} (-2)^{17} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -3 & -2 \\ 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} -1 & -2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 3 \cdot 2^{17} & 2^{18} \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} -3 \cdot 2^{17} & -2^{18} \\ 3 \cdot 2^{17} & 2^{18} \end{pmatrix}. \end{aligned}$$